

CONVERGENCE OF THE PENALTY METHOD APPLIED TO A CONSTRAINED CURVE STRAIGHTENING FLOW. *

DIETMAR OELZ †

Abstract. We apply the penalty method to the curve straightening flow of inextensible planar open curves generated by the Kirchhoff bending energy. Thus we consider the curve straightening flow of extensible planar open curves generated by a combination of the Kirchhoff bending energy and a functional penalising deviations from unit arc-length.

We start with the governing equations of the explicit parametrisation of the curve and derive an equivalent system for the two quantities indicatrix and arc-length. We prove existence and regularity of solutions and use the indicatrix/arc-length representation to compute the energy dissipation. We prove its coercivity and conclude exponential decay of the energy.

Finally, by an application of the Lions-Aubin Lemma, we prove convergence of solutions to a limit curve which is the solution of an analogous gradient flow on the manifold of inextensible open curves. This procedure also allows to characterise the Lagrange multiplier in the limit model as a weak limit of force terms present in the relaxed model.

Key words. curve straightening flow, energy dissipation, elastic regularisation, curvature flow, penalty method

subject classifications. Primary: 53C44, 35K91 Secondary: 92C10

1. Introduction

On the set of planar, open curves

$$\mathcal{A} := \{z \in H^2((0, 1), \mathbb{R}^2); b_{\min} < |z'| < b_{\max}\}$$

where $b_{\min}, b_{\max} \in \mathbb{R}$ with $0 < b_{\min} < 1 < b_{\max}$ we consider the family of gradient flows generated by the potential

$$\mathcal{E}_\varepsilon[z] := \mathcal{E}^{\text{curv}}[z] + \mathcal{E}_\varepsilon^{\text{pen}}[z]. \quad (1.1)$$

The potential consists of a functional for the curvature energy

$$\mathcal{E}^{\text{curv}}[z] := \int_0^1 \frac{|z''|^2}{2} ds \quad (1.2)$$

and of a penalising potential

$$\mathcal{E}_\varepsilon^{\text{pen}}[z] := \frac{1}{\varepsilon} \int_0^1 E(|z'|) ds, \quad (1.3)$$

where

$$E \in \mathcal{C}^2((b_{\min}, b_{\max}), \mathbb{R}_+) \quad \text{with} \quad \begin{cases} \lim_{x \rightarrow b_{\min}} E(x) = \lim_{x \rightarrow b_{\max}} E(x) = \infty, \\ E''(x) \geq \kappa > 0 \quad \forall x \in (b_{\min}, b_{\max}), \\ E'(1) = E(0) = 0. \end{cases} \quad (1.4)$$

Hence $E = E(x)$ is a strictly convex \mathcal{C}^2 -function which takes its minimum at $x = 1$. One possible choice is $E(x) = \frac{(x-1)^2}{(x-1/2)(4-x)}$. We use the functional (1.3) to penalise

*Received date / Revised version date

†Johann Radon Institute for Computational and Applied Mathematics (Linz/Vienna, Austria), (dietmar.oelz@oeaw.ac.at).

deviations from unit arc-length with a view to enforce unit arc-length by letting the scaling parameter ε tend to zero.

The gradient flow generated by the functional (1.1) is described by the system

$$\begin{cases} \partial_t z_\varepsilon + z_\varepsilon'''' - \left(\frac{1}{\varepsilon} E'(|z'_\varepsilon|) \frac{z'_\varepsilon}{|z'_\varepsilon|} \right)' = 0, \\ z_\varepsilon'', z_\varepsilon'''' - \frac{1}{\varepsilon} E'(|z'_\varepsilon|) \frac{z'_\varepsilon}{|z'_\varepsilon|} \Big|_{s=0,1} = 0, \\ z_\varepsilon(t=0, \cdot) = z_I(\cdot), \end{cases} \quad (1.5)$$

where $z_I \in \mathcal{A}$ represents the initial datum of the evolution. Here and throughout the paper $'$ denotes derivatives with respect to the arc-length. With respect to this system the centre of mass $\int_0^1 z_\varepsilon ds$ is a conserved quantity.

We show in this paper that the solution z_ε converges to a solution of the gradient flow generated by the Kirchhoff bending energy (1.2) only (cp. [Kir59, Dil92]) on the set of open, inextensible curves $\mathcal{A}_0 := \{u \in H^2((0,1), \mathbb{R}^2) : |u'| \equiv 1\}$ described by the system

$$\begin{cases} \partial_t z_0 + z_0'''' - (\lambda_0 z_0')' = 0, \\ z_0''|_{s=0,1} = 0, \\ z_0'''' - \lambda_0 z_0'|_{s=0,1} = 0, \\ |z_0'| = 1, \\ z_0(t=0, \cdot) = z_I(\cdot). \end{cases} \quad (1.6)$$

Here $\lambda_0 = \lambda_0(t, s) \in \mathbb{R}$ is a Lagrange multiplier function determined by the constraint on the arc-length

$$|z_0'| \equiv 1. \quad (1.7)$$

The system (1.5) can be seen as a regularisation of the limit system (1.6) which is treated in detail in [Oel10]. The results of this previous study are summarised in section 2. The limit model exhibits analytical properties like long time convergence at an exponential rate which, as it turns out, can be to a large extent generalised to the approximating model. As a matter of fact a large part of this study is devoted to generalising the findings on energy dissipation and large time convergence. This provides the necessary a priori bounds to prove compactness to pass to the limit as $\varepsilon \rightarrow 0$.

The system (1.5) can be used as a numerical approximation to the limit model which has appeared in the modelling of Actin-filaments in biological cells (cp. [OS10], [OS11]). Most notably in combination with an Augmented Lagrangian approach the penalising potential is currently being used in the development of numerical schemes for models in cellular biophysics.

Furthermore the results of this study provide a characterisation of the Lagrange multiplier λ_0 in the limit model (1.6) as a weak limit of a sum of forces, namely the variation of the total energy (1.1) in the direction of the arclength (compare the second line in (1.12), (4.4) and Theorem 1.5). Hence the variations of both, the curvature functional (1.2) and the penalising potential (1.3), contribute to the expression for which we show convergence to the Lagrange multiplier.

Finally it is also worth mentioning that the present study implies existence of solutions and long time convergence also for the limit problem, although these results were obtained directly in a separate study ([Oel10]).

For a short review on existing literature on curve straightening flows we refer to [Oel10]. Penalization is a popular method in optimization (e.g. [HY03]) and control theory (e.g. [CMR09]). One of the few papers we found which studies the application of the penalty method to a constrained evolution problem is [BCM10], although this is done in a totally different context.

The whole argumentation is based on an approach by which we rewrite the system (1.5) using the following notation. The symbol $b_\varepsilon = b_\varepsilon(t, s)$ denotes the arc-length

$$b_\varepsilon := |z'_\varepsilon| \quad (1.8)$$

and $\omega_\varepsilon = \omega_\varepsilon(t, s) \in \mathbb{R}$ represents the “indicatrix” of the curve z_ε (e.g. see [Lin89]) so that

$$\frac{z'_\varepsilon}{|z'_\varepsilon|} = (\cos(\omega_\varepsilon), \sin(\omega_\varepsilon)). \quad (1.9)$$

The reconstruction of the curve z_ε from the arc-length b_ε and the indicatrix ω_ε has to be done in such a way, that the centre of mass of the initial datum $\bar{z}_I := \int_0^1 z_I ds \in \mathbb{R}^2$ is conserved,

$$z_\varepsilon(t, s) = \bar{z}_I - \int_0^1 \int_0^{\bar{s}} \begin{pmatrix} \cos(\omega_\varepsilon(t, \bar{s})) \\ \sin(\omega_\varepsilon(t, \bar{s})) \end{pmatrix} b_\varepsilon d\bar{s} d\tilde{s} + \int_0^s \begin{pmatrix} \cos(\omega_\varepsilon(t, \bar{s})) \\ \sin(\omega_\varepsilon(t, \bar{s})) \end{pmatrix} d\bar{s}.$$

Observe also that $b_\varepsilon \omega'_\varepsilon$ and b'_ε are both orthogonal components of z''_ε since

$$\omega'_\varepsilon = \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|^2} \cdot z''_\varepsilon \quad \text{and} \quad b'_\varepsilon = \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot z''_\varepsilon$$

and that the total energy (1.1) can be written as

$$\mathcal{E}_\varepsilon[z] = \int_0^1 \left(\frac{1}{2} (b\omega')^2 + \frac{1}{2} (b')^2 + \frac{1}{\varepsilon} E(b) \right) ds, \quad (1.10)$$

where ω and b are the indicatrix and an arc-length of the curve z . Also the initial datum z_I will be occasionally written in terms of its indicatrix ω_I and its arc-length b_I , which allows to give two alternative expressions for the initial energy

$$\mathcal{E}_I := \int_0^1 \left[\frac{1}{2} |z_I''|^2 + \frac{1}{\varepsilon} E(|z_I'|) \right] ds = \int_0^1 \left[\frac{1}{2} (b_I \omega_I')^2 + \frac{1}{2} (b_I')^2 + \frac{1}{\varepsilon} E(b_I) \right] ds. \quad (1.11)$$

Furthermore, in view of (1.5), we use the following notation for what below will be shown to be the scalar product of $z''_\varepsilon - \frac{1}{\varepsilon} E'(|z'_\varepsilon|) \frac{z'_\varepsilon}{|z'_\varepsilon|}$ with $\frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|}$ and $\frac{z'_\varepsilon}{|z'_\varepsilon|}$ respectively, namely

$$\begin{aligned} r_\varepsilon &:= \frac{1}{b_\varepsilon} (b_\varepsilon^2 \omega'_\varepsilon)', \\ \lambda_\varepsilon &:= b_\varepsilon (\omega'_\varepsilon)^2 - b''_\varepsilon + \frac{1}{\varepsilon} E'(b_\varepsilon). \end{aligned} \quad (1.12)$$

The system (1.5) is then equivalent to

$$\begin{cases} \partial_t \omega_\varepsilon + (r''_\varepsilon - r_\varepsilon (\omega'_\varepsilon)^2 - \lambda'_\varepsilon \omega'_\varepsilon - (\lambda_\varepsilon \omega'_\varepsilon)') \frac{1}{b_\varepsilon} = 0 \\ \partial_t b_\varepsilon - r'_\varepsilon \omega'_\varepsilon - (r_\varepsilon \omega'_\varepsilon)' - \lambda''_\varepsilon + (\omega'_\varepsilon)^2 \lambda_\varepsilon = 0, \\ \omega'_\varepsilon, \omega''_\varepsilon, \lambda_\varepsilon|_{s=0,1} = 0, \\ \omega_\varepsilon(t=0, \cdot) = \omega_I(\cdot). \end{cases} \quad (1.13)$$

With respect to the energy dissipation equality

$$\frac{d}{dt}\mathcal{E}_\varepsilon = -\mathcal{D}_\varepsilon \quad (1.14)$$

this formulation allows to derive in a straightforward way that the energy dissipation is given by the two equivalent expressions

$$\mathcal{D}_\varepsilon := \int_0^1 \left[(\omega'_\varepsilon r_\varepsilon + \lambda'_\varepsilon)^2 + (r'_\varepsilon - \omega'_\varepsilon \lambda_\varepsilon)^2 \right] ds = \int_0^1 \left(z''''_\varepsilon - \left(\frac{1}{\varepsilon} E'(|z'_\varepsilon|) \frac{z'_\varepsilon}{|z'_\varepsilon|} \right)' \right)^2 ds, \quad (1.15)$$

which will be the main tool for the convergence proof. For later use we introduce a short notation for the two components of the energy dissipation,

$$m_\varepsilon := \omega'_\varepsilon r_\varepsilon + \lambda'_\varepsilon \quad \text{and} \quad n_\varepsilon := r'_\varepsilon - \omega'_\varepsilon \lambda_\varepsilon, \quad (1.16)$$

which allows to reformulate the system (1.13) as

$$\begin{cases} \partial_t \omega_\varepsilon = \frac{1}{b_\varepsilon} (\omega'_\varepsilon m_\varepsilon - n'_\varepsilon) \\ \partial_t b_\varepsilon = \omega'_\varepsilon n_\varepsilon + m'_\varepsilon, \\ \omega'_\varepsilon, \omega''_\varepsilon, \lambda_\varepsilon|_{s=0,1} = 0, \\ \omega_\varepsilon(t=0, \cdot) = \omega_I(\cdot). \end{cases} \quad (1.17)$$

In [Oel10] it was shown that the system (1.6) is equivalent to the system

$$\begin{cases} \partial_t \omega_0 + \omega_0'''' - \omega_0'^2 \omega'' - (\omega_0'' \lambda_0 + 2\omega_0' \lambda'_0) = 0, \\ -\lambda_0'' + \omega_0'^2 \lambda_0 = \omega_0'' \omega_0' + (\omega_0'' \omega_0')', \\ \omega_0', \omega_0'', \lambda_0|_{s=0,1} = 0, \\ \omega_0(t=0, \cdot) = \omega_I(\cdot). \end{cases} \quad (1.18)$$

where ω_0 represents the "indicatrix" of the curve z_0 . It was also shown in this study that the curvature energy of solutions to system (1.6) decays at an exponential rate larger or equal to $2\pi^4$ and that limit curves are straight lines.

The paper will be structured as follows. In Section 2 we summarise the results of [Oel10] as they are used in the present paper.

In Section 3 we derive the system (1.5) and prove that as a result of the usual construction in the theory of gradient flows and steepest descent flows (cp. [DGMT80], [AGS05]) there exists a weak solution z_ε to the system (1.5).

THEOREM 1.1. *Let $z_I \in \mathcal{A}$, then there is $z_\varepsilon \in H_{t,\text{loc}}^1 L_s^2 \cap C_t^{0,\frac{1}{2}} L_s^2 \cap C_t^{0,\frac{1}{8}} C_s^1 \cap L_t^\infty H_s^2$ such that z_ε is a weak solution of (1.5) satisfying*

$$\int_0^\infty \int_0^1 \left[\partial_t z_\varepsilon \cdot v + z_\varepsilon'' \cdot v'' + \frac{1}{\varepsilon} E'(|z'_\varepsilon|) \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot v' \right] ds dt = 0 \quad (1.19)$$

for all $v \in C_c^\infty(\mathbb{R}_+, C_s^\infty)$ and $z_\varepsilon(t=0, \cdot) = z_I$. It holds that $\partial_t z_\varepsilon \in L_t^2 L_s^2$ with $\|\partial_t z_\varepsilon\|_{L_t^2 L_s^2} \leq \sqrt{2\mathcal{E}_I}$.

Here and in the sequel we abbreviate the notation of function spaces writing the subscripts

$$\begin{aligned} \square_t & \text{ for function spaces on } \{t \in [0, \infty)\} \quad \text{and} \\ \square_s & \text{ for function spaces on } \{s \in [0, 1]\}. \end{aligned}$$

In Section 4 we rewrite (1.5) obtaining the system (1.13) and prove

THEOREM 1.2. *The solution according to Theorem 1.1 gives a distributional sense to the system (1.13), i.e. for ω_ε being the indicatrix of z_ε and b_ε its arc-length and using the notation (1.12) it holds that*

$$\int_0^\infty \int_0^1 [-b_\varepsilon \phi_t + (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) \phi' + (\omega'_\varepsilon \lambda_\varepsilon - r'_\varepsilon) \omega'_\varepsilon \phi] ds dt = 0, \quad (1.20)$$

$$\int_0^\infty \int_0^1 \left[-\omega_\varepsilon \partial_t \psi - (r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon) \left(\frac{\psi}{b_\varepsilon} \right)' - (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) \frac{\omega'_\varepsilon}{b_\varepsilon} \psi \right] ds dt = 0. \quad (1.21)$$

for all $\psi, \phi \in H_{0,t}^1 L_s^2 \cap L_t^2 H_{0,s}^1$ and $\lambda_\varepsilon, \omega'_\varepsilon, \omega''_\varepsilon|_{0,1} = 0$ a.e. on \mathbb{R}_+ . Furthermore it holds that $\omega_\varepsilon, b_\varepsilon \in \mathcal{C}_t^{0,1/8} \mathcal{C}_s^0 \cap L_t^\infty H_s^1$ and $\lambda_\varepsilon, r_\varepsilon \in L_t^2 H_s^1$. For every $T > 0$ it also holds that $\omega'''_\varepsilon, b''_\varepsilon \in L^2(0, T; L_s^1)$ uniformly with respect to ε .

Next, in Section 5, we show the formal derivation of the energy dissipation (1.15) and show that the energy dissipation equation (1.14) is satisfied in a weak sense.

THEOREM 1.3. *Let $z_I \in \mathcal{A}$, let z_ε be a solution of problem (1.5) according to Theorem 1.1 and let $(\omega_\varepsilon, b_\varepsilon, \lambda_\varepsilon)$ be the corresponding solution to (1.13) according to Theorem 1.2, then the energy dissipation equality (1.14) holds weakly in time.*

Finally in Section 6 we prove coercivity of (1.15) with respect to the total energy given by (1.1) and (1.10) (a Poincaré type inequality) obtaining the exponential decay of the energy.

THEOREM 1.4. *(Poincaré-type) Under the assumptions of Theorem 1.3 there is a constant $C > 0$ such that*

$$\mathcal{E}_\varepsilon \leq \mathcal{E}_I \exp(-Ct).$$

Most notably at large times the total energy tends to zero, $\lim_{t \rightarrow \infty} \mathcal{E}_\varepsilon = 0$.

Finally we obtain the main theorem of this paper in Section 7. It states the convergence as $\varepsilon \rightarrow 0$ and the consistency with the limit system, i.e. a subsequence of solutions to the problem (1.13) and (1.5) converges to a solution of (1.18) and (1.6) respectively.

THEOREM 1.5. *Let $(\omega_\varepsilon, b_\varepsilon, \lambda_\varepsilon)$ as in Theorem 1.2 be a solution to the system (1.13), then there is a subsequence $\varepsilon_i \rightarrow 0$ and limit functions (ω_0, λ_0) such that*

$$\begin{aligned} b_{\varepsilon_i} &\rightarrow \mathbf{1} && \text{in } L^1((0, T), W_s^{1,r}) \cap L^q((0, T), H_s^1) \quad (1 \leq r, q < \infty), \\ \omega_{\varepsilon_i} &\rightarrow \omega_0 && \text{in } L^1((0, T), W_s^{2,r}) \cap L^q((0, T), H_s^1) \quad (1 \leq r, q < \infty), \\ m_{\varepsilon_i} &\rightarrow \omega'_0 \omega''_0 + \lambda'_0 && \text{in } L_t^2 L_s^2, \\ n_{\varepsilon_i} &\rightarrow \omega'''_0 - \omega'_0 \lambda_0 && \text{in } L_t^2 L_s^2, \\ r_{\varepsilon_i} &\rightarrow \omega''_0 && \text{in } L^1((0, T), L_s^2) \quad \text{and} \quad r'_{\varepsilon_i} \rightharpoonup \omega'''_0 && \text{in } L_t^2 L_s^2, \\ \lambda_{\varepsilon_i} &\rightharpoonup \lambda_0 && \text{in } L_t^2 H_s^1, \end{aligned}$$

as $\varepsilon_i \rightarrow 0$, where we use the following notation for constant functions, $\mathbf{1}: [0, \infty) \times [0, 1] \mapsto \mathbb{R}$, $x \mapsto 1$ and $\mathbf{0}: [0, \infty) \times [0, 1] \mapsto \mathbb{R}$, $x \mapsto 0$.

The limit functions ω_0, λ_0 satisfy the weak formulation of (1.18) and they also allow to reconstruct z_0 which is then a weak solution of (1.6).

2. Preliminary and technical results We cite here some of the results we obtained in [Oel10] for the limit system (1.6) and the equivalent system satisfied by the indicatrix (1.18). We define the constrained set of curves

$$\mathcal{A}_0 := \{z \in H^2([0, 1], \mathbb{R}^2) : |z'| \equiv 1\}.$$

THEOREM 2.1. Let $z_I \in \mathcal{A}_0$, then there are $z_0 \in H_{t,\text{loc}}^1 L_s^2 \cap C_t^{0,\frac{1}{2}} L_s^2 \cap C_t^{0,\frac{1}{8}} C_s^1 \cap L_t^\infty H_s^2$ and $\bar{\lambda}_0 \in L_{\text{loc},t}^2 \mathcal{M}_s$ such that the pair $(z_0, \bar{\lambda}_0)$ is a weak solution of (1.6) satisfying

$$\int_0^\infty \int_0^1 [z_0'' \cdot v'' + \partial_t z_0 \cdot v + \bar{\lambda}_0 z_0' \cdot v'] ds dt = 0 \quad (2.1)$$

for all $v \in C_c^\infty(\mathbb{R}_+, C_s^\infty)$, $z_0(t=0, \cdot) = z_{0,I}$ and the constraint (1.7) in a pointwise sense for all $t \geq 0$ and $s \in [0, 1]$.

THEOREM 2.2. The solution according to Theorem 2.1 gives a distributional sense to the system (1.18), i.e. for ω_0 being the indicatrix of z_0 and $\lambda_0 = \bar{\lambda}_0 + \omega'^2$ it holds that

$$\int_0^\infty \int_0^1 [-\omega_0 \partial_t \psi - \omega_0''' \psi' - \omega_0'' (\omega_0')^2 \psi - \lambda_0' \omega_0' \psi + \lambda_0 \omega_0' \psi'] ds dt = 0 \quad \text{and} \quad (2.2)$$

$$\int_0^\infty \int_0^1 [-\omega_0''' \omega_0' \phi + \omega_0'' \omega_0' \phi' + \lambda_0' \phi' + \lambda_0 (\omega_0')^2 \phi] ds dt = 0. \quad (2.3)$$

for all $\psi \in H_{0,t}^1 L_s^2 \cap L_t^2 H_{0,s}^1$ and $\phi \in L_t^2 H_{0,s}^1$ and $\lambda_0, \omega_0', \omega_0''|_{0,1} = 0$ a.e. on \mathbb{R}_+ . Furthermore it holds that $\omega_0 \in C_t^{0,1/8} C_s^0$ with $\omega_0' \in L_t^\infty L_s^2 \cap L_t^2 H_s^2$ and $\lambda_0 \in L_t^2 H_s^1$.

THEOREM 2.3. Let $z_I \in \mathcal{A}_0$, let $(z_0, \bar{\lambda}_0)$ be a solution of problem (1.6) according to Theorem 2.1 and let (ω_0, λ_0) be the corresponding solution to (1.18) according to Theorem 2.2. The curvature energy can then be equivalently formulated in terms of z_0 and in terms of ω_0 as

$$\mathcal{E}_0 := \int_0^1 \frac{1}{2} |z_0''|^2 ds = \int_0^1 \frac{1}{2} (\omega_0')^2 ds \quad (2.4)$$

and the curvature energy of the initial datum is given by

$$\mathcal{E}_{0,I} := \int_0^1 \frac{1}{2} |z_{0,I}''|^2 ds. \quad (2.5)$$

The energy dissipation is then given by

$$\frac{d}{dt} \mathcal{E}_0 = -\mathcal{D}_0 \quad \text{with} \quad \mathcal{D}_0 := \int_0^1 [(\omega_0' \omega_0'' + \lambda_0')^2 + (\omega_0''' - \omega_0' \lambda_0)^2] ds, \quad (2.6)$$

in the sense that (2.6) holds weakly in time.

Finally the coercivity of \mathcal{D}_0 with respect to the curvature energy \mathcal{E}_0 (a Poincaré type inequality) yields the exponential decay of the energy.

THEOREM 2.4. (Poincaré-type) Under the assumptions of Theorem 2.3, let the energy of the initial datum be given by (2.5), then it holds that

$$\mathcal{E}_0 \leq \mathcal{E}_{0,I} \exp(-2\pi^4 t),$$

where again the curvature energy is alternatively given by (2.4).

As a consequence it holds that

$$\|\partial_t z_0\|_{L_t^2 L_s^2}^2 = \int_0^\infty \mathcal{D}_0 dt = \mathcal{E}_{0,I}. \quad (2.7)$$

Finally we add here a technical statement which derives from the definition of the integrand of the punishing potential (1.4) two inequalities that we will use below in Section 6 and Section 7.

LEMMA 2.5. *Let the punishing profile be as defined in (1.4), which implies among other things that $E''(x) \geq \kappa$ for all $x \in (b_{\min}, b_{\max})$, then it satisfies*

$$\frac{\kappa}{2}(x-1)^2 \leq E(x) \leq \frac{1}{2\kappa}E'(x)^2$$

for all $x \in (b_{\min}, b_{\max})$

Proof. If $1 \leq x < b_{\max}$ integrate $E''(x) \geq \kappa$ twice on $(1, x)$ to obtain the first inequality. For the second inequality use $E'(x) \geq 0$ to obtain $E'(x)E''(x) \geq \kappa E'(x)$ which gives the result after one integration.

If $b_{\min} < x < 1$ integrate instead on $(x, 1)$ and use $-E'(x) \geq 0$. \square

3. Existence of solutions The definition of the curvature energy (1.2) and the notation used in (1.10) imply that

$$\int_0^1 (\omega')^2 ds \leq 2b_{\min}^{-2} \mathcal{E}^{\text{curv}}[z], \quad \int_0^1 (b')^2 ds \leq 2 \mathcal{E}^{\text{curv}}[z] \quad \text{and} \quad \int_0^1 |z''|^2 ds \leq 2 \mathcal{E}^{\text{curv}}[z]. \quad (3.1)$$

We introduce the time step approximation scheme

$$Z_\varepsilon^0 = z_I \quad \text{and} \quad Z_\varepsilon^n \in \operatorname{argmin}_{w: [0,1] \rightarrow \mathbb{R}} \{ \mathcal{E}_\tau(Z_\varepsilon^{n-1})[w] + \mathcal{E}_\varepsilon[w] \}, \quad (3.2)$$

where $\tau > 0$ is the constant size of the time steps, $n = 0, 1, \dots$ the index of the respective time step and

$$\mathcal{E}_\tau(Z_\varepsilon^{n-1})[w] := \int_0^1 \frac{|w(s) - Z_\varepsilon^{n-1}|^2}{2\tau} ds.$$

LEMMA 3.1. *For any given $u \in \mathcal{A}$ there are constants $C_1 > 0$ and $C_2 > 0$ such that*

$$\mathcal{E}_\tau(u)[w] + \mathcal{E}_\varepsilon[w] \geq C_1 \|w\|_{H_s^2} - C_2.$$

Proof. Here we use (3.1) and the fact that the norm $\|w\| := \|w\|_{L^2} + \|\partial_{ss} w\|_{L^2}$ is equivalent to the usual norm in H_s^2 . \square

LEMMA 3.2. *Let $\tau > 0$ and $Z_\varepsilon^n \in \mathcal{A}$. Then $\mathcal{E}^{\text{curv}}$ is weakly lower semicontinuous and $\mathcal{E}_\varepsilon^{\text{pen}}$ and \mathcal{E}_τ are weakly continuous with respect to the $H^2(0,1)^2$ -topology.*

Proof. Weak lower semicontinuity is a consequence of the convexity of $\mathcal{E}^{\text{curv}}$. The integrand of \mathcal{E}_τ only depends on w in a Lipschitz continuous way. Since due to Lemma 3.1 and embedding into C^1 there is a $\delta > 0$ such that $b_{\min} + \delta \leq |w'| \leq b_{\max} - \delta$ the same is true for $\mathcal{E}_\varepsilon^{\text{pen}}$ with respect to w' . The result is therefore a consequence of the compact embedding of $H^2(0,1)$ in $C^1([0,1])$. \square This proves the existence of a sequence $(Z_\varepsilon^n)_{n=0,1,\dots}$ defined by (3.2) with $Z_\varepsilon^n \in H_s^2$. Approximations of the solution of the continuous problem are then defined by linear interpolation and by piecewise constant extension:

$$\left. \begin{aligned} Z_{\varepsilon,\tau}(t,s) &:= Z_\varepsilon^n(s) + \left(\frac{t}{\tau} - n\right) (Z_\varepsilon^{n+1}(s) - Z_\varepsilon^n(s)), \\ Z_{\varepsilon,\tau}^{\text{old}}(t,s) &:= Z_\varepsilon^n(s), \\ Z_{\varepsilon,\tau}^{\text{new}}(t,s) &:= Z_\varepsilon^{n+1}(s), \end{aligned} \right\} \quad \text{for } n\tau < t \leq (n+1)\tau.$$

LEMMA 3.3. For every fixed finite $T > 0$, $Z_{\varepsilon, \tau} \in H^1((0, T), L^2(0, 1))$ uniformly in τ .

Proof. The variational principle (3.2) implies that

$$\int_0^1 \left[\frac{1}{2\tau} (Z_\varepsilon^n - Z_\varepsilon^{n-1})^2 \right] ds + \mathcal{E}_\varepsilon[Z_\varepsilon^n] \leq \mathcal{E}_\varepsilon[Z_\varepsilon^{n-1}]. \quad (3.3)$$

As a consequence it holds that

$$\frac{1}{2\tau} \|Z_\varepsilon^{n+1} - Z_\varepsilon^n\|_{L^2(0,1)}^2 \leq \mathcal{E}_\varepsilon[Z_\varepsilon^n] - \mathcal{E}_\varepsilon[Z_\varepsilon^{n+1}]. \quad (3.4)$$

Since the time derivative of $Z_{\varepsilon, \tau}$ is piecewise constant taking the sum $n = 0, 1, \dots, m-1$ in (3.4) where $m = \lceil T/\tau \rceil$ implies

$$\int_0^{m\tau} \|\partial_t Z_{\varepsilon, \tau}\|_{L^2(0,1)}^2 dt = \frac{1}{\tau} \sum_{n=0}^{m-1} \|Z_\varepsilon^{n+1} - Z_\varepsilon^n\|_{L^2(0,1)}^2 \leq 2\mathcal{E}_\varepsilon[z_I], \quad (3.5)$$

completing the proof. \square

This result sets the stage for passing to the limit in the approximate solutions.

LEMMA 3.4. For every fixed finite $T > 0$,

$$\lim_{\tau \rightarrow 0} Z_{\varepsilon, \tau} = z_\varepsilon \in L^\infty((0, T); H^2(0, 1)) \cap C^{0,1/8}([0, T]; C^1([0, 1])) \cap H^1((0, T); L^2(0, 1)),$$

(restricting to subsequences) where the convergence is strong in $C([0, T]; C^1([0, 1]))$, weak in $H^1((0, T); L^2(0, 1))$, and weak* in $L^\infty((0, T); H^2(0, 1))$. The piecewise constant approximations $Z_{\varepsilon, \tau}^{\text{old}}$ and $Z_{\varepsilon, \tau}^{\text{new}}$ converge to z strongly in $L^\infty([0, T]; C^1([0, 1]))$ and weakly* in $L^\infty((0, T); H^2(0, 1))$.

Proof. The inequalities (3.3) and (3.1) imply that $(Z_{\varepsilon, \tau})_\tau$ is bounded in $L^\infty((0, T); H^2(0, 1))$ uniformly with respect to τ , which already shows the weak* convergence. The weak convergence is a consequence of the previous lemma since the inequality (3.5) for $T \rightarrow \infty$ implies that

$$\|\partial_t z_\varepsilon\|_{L_t^2 L_s^2}^2 \leq 2\mathcal{E}_\varepsilon[z_I]. \quad (3.6)$$

Another consequence is that $Z_{\varepsilon, \tau}$ is uniformly bounded in $C^{0,1/2}([0, T]; L^2(0, 1))$. The interpolation inequality

$$\|u\|_{C^{1,\alpha}([0,1])} \leq c \|u\|_{L^2(0,1)}^{(1-2\alpha)/4} \|u\|_{H^2(0,1)}^{(3+2\alpha)/4}$$

for $0 \leq \alpha \leq 1/2$, can then be used together with the $H^2(0,1)$ -bound to obtain

$$\|Z_{\varepsilon, \tau}(t_2) - Z_{\varepsilon, \tau}(t_1)\|_{C^{1,\alpha}([0,1])} \leq c_T |t_2 - t_1|^{(1-2\alpha)/8},$$

completing the convergence proof for $Z_{\varepsilon, \tau}$ by an application of the Arzela-Ascoli theorem. The convergence results for $Z_{\varepsilon, \tau}^{\text{old}}$ and $Z_{\varepsilon, \tau}^{\text{new}}$ are straightforward consequences. \square

This implies the Theorem 1.1 due to the following argumentation.

Proof. of Theorem 1.1. By construction the variational equation of (3.2)

$$\delta \mathcal{E}_\tau(Z_\varepsilon^n)[Z_\varepsilon^{n+1}]v(t, \cdot) + \delta \mathcal{E}_\varepsilon(Z_\varepsilon^n)[Z_\varepsilon^{n+1}]v(t, \cdot) = 0,$$

holds for a test function $v \in \mathcal{C}_c^\infty(\mathbb{R}_+, \mathcal{C}_s^\infty)$ and $n\tau < t \leq (n+1)\tau$. With the definitions of $Z_{\varepsilon, \tau}$, $Z_{\varepsilon, \tau}^{\text{old}}$, and $Z_{\varepsilon, \tau}^{\text{new}}$ this can be written as

$$\int_0^1 \left[\partial_t Z_{\varepsilon, \tau} \cdot v + \partial_s^2 Z_{\varepsilon, \tau}^{\text{new}} \cdot \partial_s^2 v + \frac{1}{\varepsilon} E'(|\partial_s Z_{\varepsilon, \tau}^{\text{new}}|) \frac{\partial_s Z_{\varepsilon, \tau}^{\text{new}}}{|\partial_s Z_{\varepsilon, \tau}^{\text{new}}|} \cdot \partial_s v \right] ds = 0.$$

After integration with respect to t , we pass to the limit. Note that the weakly convergent terms $\partial_s^2 Z_{\varepsilon, \tau}^{\text{new}}$ and $\partial_t Z_{\varepsilon, \tau}$ occur only linearly, and that all other terms converge strongly. \square

4. Indicatrix representation As mentioned above, r_ε and λ_ε , which are defined in (1.12), are two components of $z_\varepsilon''' - \frac{1}{\varepsilon} E'(|z_\varepsilon'|) \frac{z_\varepsilon'}{|z_\varepsilon'|}$, namely

$$r_\varepsilon = \left(z_\varepsilon''' - \frac{1}{\varepsilon} E'(|z_\varepsilon'|) \frac{z_\varepsilon'}{|z_\varepsilon'|} \right) \cdot \frac{z_\varepsilon'^\perp}{|z_\varepsilon'|} = z_\varepsilon''' \cdot \frac{z_\varepsilon'^\perp}{|z_\varepsilon'|}, \quad (4.1)$$

$$\lambda_\varepsilon = \left(z_\varepsilon''' - \frac{1}{\varepsilon} E'(|z_\varepsilon'|) \frac{z_\varepsilon'}{|z_\varepsilon'|} \right) \cdot \frac{z_\varepsilon'}{|z_\varepsilon'|} = z_\varepsilon''' \cdot \frac{z_\varepsilon'}{|z_\varepsilon'|} - \frac{1}{\varepsilon} E'(|z_\varepsilon'|). \quad (4.2)$$

Hence the system (1.5) can be written as

$$\begin{cases} \partial_t z_\varepsilon + \left(r_\varepsilon \frac{z_\varepsilon'^\perp}{|z_\varepsilon'|} \right)' - \left(\lambda_\varepsilon \frac{z_\varepsilon'}{|z_\varepsilon'|} \right)' = 0, \\ z_\varepsilon'', r_\varepsilon, \lambda_\varepsilon \Big|_{s=0,1} = 0, \\ z_\varepsilon(t=0, \cdot) = z_I(\cdot). \end{cases} \quad (4.3)$$

The definitions in (1.12) are motivated by the variation of \mathcal{E}_ε as formulated in (1.10),

$$\begin{aligned} \delta \mathcal{E}_\varepsilon[\omega, b](\delta\omega, \delta b) &= \int_0^1 \left(b^2 \omega' (\delta\omega)' + b (\omega')^2 \delta b + b' \delta b' + \frac{1}{\varepsilon} E'(b) \delta b \right) ds = \\ &= [b^2 \omega' \delta\omega + b' \delta b]_0^1 + \int_0^1 - (b^2 \omega')' \delta\omega + \left(b (\omega')^2 - b'' + \frac{1}{\varepsilon} E'(b) \right) \delta b ds. \end{aligned}$$

Replacing the variations of the indicatrix and of the arc-length by $\delta\omega = z'^\perp \cdot \delta z' / |z'|^2$ and $\delta b = z' \cdot \delta z' / |z'|$ respectively we obtain

$$\begin{aligned} \delta \mathcal{E}_\varepsilon \delta z &= \left[\left(b \omega' \frac{z'^\perp}{|z'|} + b' \frac{z'}{|z'|} \right) \cdot \delta z' \right]_0^1 + \\ &+ \int_0^1 \delta z' \cdot \left(-\frac{1}{b} (b^2 \omega')' \frac{z'^\perp}{|z'|} + \left(b (\omega')^2 - b'' + \frac{1}{\varepsilon} E'(b) \right) \frac{z'}{|z'|} \right) ds, \quad (4.4) \end{aligned}$$

which motivates the definitions (1.12). The boundary terms in (4.4) also suggest to replace the boundary condition $z_\varepsilon''|_{s=0,1} = 0$ by $\omega'_\varepsilon, b'_\varepsilon|_{s=0,1} = 0$.

LEMMA 4.1. *Let $z_I \in \mathcal{A}$, let z_ε be a weak solution to problem (1.5) according to Theorem 1.1, then there is $\omega_\varepsilon \in \mathcal{C}_t^{0,1/8} \mathcal{C}_s^0 \cap L_t^\infty H_s^1$ such that ω_ε is an indicatrix of z_ε . Furthermore it holds that $b_\varepsilon \in \mathcal{C}_t^{0,1/8} \mathcal{C}_s^0 \cap L_t^\infty H_s^1$ and $(z_\varepsilon, \omega_\varepsilon, b_\varepsilon)$ constitute a weak solution of (4.3) satisfying*

$$\begin{aligned} \int_0^\infty \int_0^1 \left[\partial_t z_\varepsilon \cdot v + b_\varepsilon^2 \omega'_\varepsilon \left(\frac{1}{b_\varepsilon} \frac{z_\varepsilon'^\perp}{|z_\varepsilon'|} \cdot v'_\varepsilon \right)' + \right. \\ \left. + \left(b_\varepsilon (\omega'_\varepsilon)^2 + \frac{1}{\varepsilon} E'(b_\varepsilon) \right) \frac{z_\varepsilon'}{|z_\varepsilon'|} \cdot v' + b'_\varepsilon \left(\frac{z_\varepsilon'}{|z_\varepsilon'|} \cdot v' \right)' \right] ds dt = 0 \quad (4.5) \end{aligned}$$

for all $v \in C_c^\infty(\mathbb{R}_+, C_s^\infty)$.

Proof. The regularity of z'_ε according to Theorem 1.1 allows to identify $\omega_\varepsilon \in C_t^{0,1/8} C_s^0 \cap L_t^\infty H_s^1$ up to an additive constant being a multiple of 2π . The regularity of b_ε is due to its definition and the result of Theorem 1.1. The rest of the statement is then a consequence of the discussion above. \square

LEMMA 4.2. (*Regularity, a-priori estimates*) Let $(z_\varepsilon, \omega_\varepsilon, \lambda_\varepsilon)$ bet a weak solution of (4.3) in the sense of Lemma 4.1, where we use the notation (1.12), then it holds that $r_\varepsilon \in L_t^2 L_s^\infty$, $\omega'_\varepsilon r_\varepsilon \in L_t^2 L_s^2$, $\lambda_\varepsilon \in L_t^2 L_s^\infty$, $\omega'_\varepsilon \lambda_\varepsilon \in L_t^2 L_s^2$, $r'_\varepsilon \in L_t^2 L_s^2$ and $\lambda'_\varepsilon \in L_t^2 L_s^2$. It also holds that

$$\int_0^\infty \int_0^1 \left[(r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon)^2 + (\lambda'_\varepsilon + \omega'_\varepsilon r_\varepsilon)^2 \right] ds dt = \|\partial_t z_\varepsilon\|_{L_t^2 L_s^2}^2.$$

Proof. First note that $\int_0^s \partial_t z_\varepsilon \in L_t^2 L_s^\infty$ since

$$\int_0^\infty \left\| \int_0^s \partial_t z_\varepsilon ds \right\|_\infty^2 dt \leq \int_0^\infty \left(\int_0^1 |\partial_t z_\varepsilon| ds \right)^2 dt \leq \int_0^\infty \int_0^1 |\partial_t z_\varepsilon|^2 ds dt < 2\mathcal{E}_I$$

by (3.6). We go back to the integrated version of (4.3)

$$\begin{cases} \int_0^s \partial_t z_\varepsilon ds + r_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} - \lambda_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} = 0, \\ \omega'_\varepsilon, b'_\varepsilon \Big|_{s=0,1} = 0, \\ z_\varepsilon(t=0, \cdot) = z_I(\cdot), \end{cases}$$

where $\omega'_\varepsilon, b'_\varepsilon \Big|_{s=0,1} = 0$ is an equivalent notation for $z''_\varepsilon \Big|_{s=0,1} = 0$ and obtain

$$\frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot \int_0^s \partial_t z_\varepsilon ds = -r_\varepsilon \quad \text{and} \quad \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot \int_0^s \partial_t z_\varepsilon ds = \lambda_\varepsilon,$$

which implies $r_\varepsilon \in L_t^2 L_s^\infty$ and $\lambda_\varepsilon \in L_t^2 L_s^\infty$.

Moreover we find that

$$\begin{aligned} \int_0^\infty \int_0^1 (\lambda_\varepsilon \omega'_\varepsilon)^2 ds dt &\leq \int_0^\infty \|\lambda_\varepsilon\|_{L_s^\infty}^2 \int_0^1 (\omega'_\varepsilon)^2 ds dt \leq \\ &\leq 2\mathcal{E}_I b_{\min}^{-2} \int_0^\infty \|\lambda_\varepsilon\|_{L_s^\infty}^2 dt < (2\mathcal{E}_I)^2 b_{\min}^{-2}, \end{aligned} \quad (4.6)$$

where we used (3.1). This computation and an analogous one for $\omega'_\varepsilon r_\varepsilon$ imply

$$\omega'_\varepsilon \lambda_\varepsilon \in L_t^2 L_s^2 \quad \text{and} \quad \omega'_\varepsilon r_\varepsilon \in L_t^2 L_s^2. \quad (4.7)$$

We write the weak formulation (4.5) of problem (4.3) after two integrations by parts,

$$\begin{aligned} \int_0^\infty \int_0^1 \left(\partial_t z_\varepsilon + \left(r_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \right)' - \left(\lambda_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \right)' \right) \cdot v ds dt + \\ + \int_0^\infty \left[-r_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot v + \lambda_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot v + b_\varepsilon \omega'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot v' + b'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot v' \right]_0^1 dt = 0, \end{aligned} \quad (4.8)$$

for all $v \in C_c^\infty(\mathbb{R}_+, C_s^\infty)$.

The uniform estimates we obtained allow to set $v = \frac{z_\varepsilon^\perp}{|z_\varepsilon^\perp|} \phi$ for a test function $\phi \in \mathcal{D}([0, \infty) \times [0, 1])$ and to obtain $\int_0^\infty \int_0^1 \left[\partial_t z_\varepsilon \cdot \frac{z_\varepsilon^\perp}{|z_\varepsilon^\perp|} \phi + r'_\varepsilon \phi - \lambda_\varepsilon \omega'_\varepsilon \phi \right] ds dt = 0$.

Specialising (4.8) for $v = \frac{z'_\varepsilon}{|z'_\varepsilon|} \psi$ with $\psi \in \mathcal{D}([0, \infty) \times [0, 1])$ we conclude by an analogous computation $\int_0^\infty \int_0^1 \left[\partial_t z_\varepsilon \cdot \frac{z'_\varepsilon}{|z'_\varepsilon|} \psi - \omega'_\varepsilon r_\varepsilon \psi - \lambda'_\varepsilon \psi \right] ds dt = 0$. Due to (3.6) and (4.7) this implies

$$\frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot \partial_t z_\varepsilon + r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon = 0 \quad \text{a.e.} \quad \text{and} \quad \frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot \partial_t z_\varepsilon - \omega'_\varepsilon r_\varepsilon - \lambda'_\varepsilon = 0 \quad \text{a.e.}, \quad (4.9)$$

hence $r'_\varepsilon \in L_t^2 L_s^2$ and $\lambda'_\varepsilon \in L_t^2 L_s^2$. Finally (4.9) implies

$$\begin{aligned} \int_0^\infty \int_0^1 \left[(r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon)^2 + (\lambda'_\varepsilon + \omega'_\varepsilon r_\varepsilon)^2 \right] ds dt &= \\ &= \int_0^\infty \int_0^1 \left[\left(\frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot \partial_t z_\varepsilon \right)^2 + \left(\frac{z'_\varepsilon}{|z'_\varepsilon|} \cdot \partial_t z_\varepsilon \right)^2 \right] ds dt = \|\partial_t z_\varepsilon\|_{L_t^2 L_s^2}^2. \end{aligned} \quad (4.10)$$

□

Using this result we also get insight into regularity of higher derivatives of ω_ε and b_ε .

LEMMA 4.3. (*Regularity of higher derivatives*) *Let $(z_\varepsilon, \omega_\varepsilon, \lambda_\varepsilon)$ be a weak solution of (4.3) in the sense of Lemma 4.1, then it holds for every $T > 0$ that the family $\{\omega_\varepsilon\}$ is uniformly bounded in $L^1(0, T; W^{3,1}(0, 1))$ and that the family $\{b_\varepsilon\}$ is uniformly bounded in $L^1(0, T; W^{2,1}(0, 1))$. It also holds that $b_\varepsilon''' \in L^2(0, T; L_s^1)$, but not uniformly with respect to ε .*

Proof. For the proof we use the definitions in (1.12). It holds that $\omega_\varepsilon'' = \frac{r_\varepsilon - 2b'_\varepsilon \omega'_\varepsilon}{b_\varepsilon}$, hence

$$\|\omega_\varepsilon''\|_{L^2(0, T; L_s^1)} \leq \frac{1}{b_{\min}} \left(\|r_\varepsilon\|_{L^2(0, T; L_s^2)} + 2\|b'_\varepsilon\|_{L^2(0, T; L_s^2)} \|\omega'_\varepsilon\|_{L^2(0, T; L_s^2)} \right),$$

which is uniformly bounded due to Lemma 4.2 and Lemma 4.1.

Furthermore the definition of r_ε in (1.12) also implies that $\omega_\varepsilon''' = r'_\varepsilon - 2b''_\varepsilon \omega'_\varepsilon - 3b'_\varepsilon \omega_\varepsilon'' / b_\varepsilon$. Hence we estimate

$$\begin{aligned} \|\omega_\varepsilon'''\|_{L^1(0, T; L_s^1)} &\leq b_{\min}^{-1} \left(\|r'_\varepsilon\|_{L^1(0, T; L_s^1)} + 2\|b''_\varepsilon\|_{L^2(0, T; L_s^1)} \|\omega'_\varepsilon\|_{L^2(0, T; L_s^\infty)} + \right. \\ &\quad \left. + 3\|b'_\varepsilon\|_{L^2(0, T; L_s^\infty)} \|\omega_\varepsilon''\|_{L^2(0, T; L_s^1)} \right) \\ &\leq b_{\min}^{-1} \left(\sqrt{T} \|r'_\varepsilon\|_{L^2(0, T; L_s^2)} + 5\|b''_\varepsilon\|_{L^2(0, T; L_s^1)} \|\omega_\varepsilon''\|_{L^2(0, T; L_s^1)} \right) \end{aligned}$$

which is uniformly bounded due to the result above.

Observe that due to Lemma 3.1 and embedding into C^1 there is a $\delta > 0$ such that $b_{\min} + \delta \leq |z'_\varepsilon| \leq b_{\max} - \delta$. Hence, using $b''_\varepsilon = b_\varepsilon (\omega'_\varepsilon)^2 - \lambda_\varepsilon + \frac{1}{\varepsilon} E'(b_\varepsilon)$ we conclude that

$$\|b''_\varepsilon\|_{L^2(0, T; L_s^1)} \leq b_{\max} \|\omega'_\varepsilon\|_{L^2(0, T; L_s^2)} + 2\|\lambda_\varepsilon\|_{L^2(0, T; L_s^2)} + \frac{1}{\varepsilon} \max_{b_{\min} + \delta \leq x \leq b_{\max} - \delta} E'(x), \quad (4.11)$$

is bounded due to Lemma 4.2 and Lemma 4.1. This bound is not uniform with respect to ε , but can be replaced by a uniform one using the following computation.

Integrating the definition of λ_ε against $E'(b_\varepsilon)$ we obtain

$$\int_0^1 \lambda_\varepsilon E'(b_\varepsilon) d\tilde{s} = \int_0^1 \left[b_\varepsilon (\omega'_\varepsilon)^2 E'(b_\varepsilon) + b_\varepsilon'^2 E''(b_\varepsilon) + \frac{1}{\varepsilon} E'(b_\varepsilon)^2 \right] d\tilde{s}.$$

This implies the inequality

$$\int_0^1 \frac{1}{\varepsilon} E'(b_\varepsilon)^2 d\tilde{s} \leq \int_0^1 (\lambda_\varepsilon E'(b_\varepsilon) - b_\varepsilon (\omega'_\varepsilon)^2 E'(b_\varepsilon)) d\tilde{s} \leq \|\lambda_\varepsilon - b_\varepsilon (\omega'_\varepsilon)^2\|_{L^2} \|E'(b_\varepsilon)\|_{L^2}.$$

Hence we conclude that

$$\frac{1}{\varepsilon} \|E'(b_\varepsilon)\|_{L^2} \leq \|\lambda_\varepsilon - b_\varepsilon (\omega'_\varepsilon)^2\|_{L^2}, \quad (4.12)$$

which allows to replace (4.11) by

$$\|b_\varepsilon''\|_{L^2(0,T;L_s^1)} \leq b_{\max} \|\omega'_\varepsilon\|_{L^2(0,T;L_s^2)} + 2\|\lambda_\varepsilon\|_{L^2(0,T;L_s^2)} + \|\lambda_\varepsilon - b_\varepsilon (\omega'_\varepsilon)^2\|_{L^2(0,T;L_s^2)},$$

which is uniformly bounded due to the results of Lemma 4.2 and Lemma 4.1.

Furthermore the definition of λ_ε in (1.12) also implies that $b_\varepsilon''' = b_\varepsilon' (\omega'_\varepsilon)^2 + 2b_\varepsilon \omega'_\varepsilon \omega''_\varepsilon - \lambda'_\varepsilon + \frac{1}{\varepsilon} E''(b_\varepsilon) b'_\varepsilon$. Hence we estimate

$$\begin{aligned} \|b_\varepsilon'''\|_{L^1(0,T;L_s^1)} &\leq \|b'_\varepsilon\|_{L^1(0,T;L_s^\infty)} \|(\omega'_\varepsilon)^2\|_{L^\infty(0,T;L_s^1)} + 2b_{\max} \|\omega''_\varepsilon\|_{L^2(0,T;L_s^1)}^2 + \\ &\quad + \|\lambda'_\varepsilon\|_{L^1(0,T;L_s^2)} + \frac{1}{\varepsilon} \|b'_\varepsilon\|_{L^1(0,T;L_s^1)} \max_{b_{\min} + \delta \leq x \leq b_{\max} - \delta} E''(x) \end{aligned}$$

which is bounded due to the result above, but not uniformly with respect to ε .

□

Next we will derive the governing equations for the quantities ω_ε and b_ε making use of the following identities

$$\left(\frac{z'}{|z'|} \right)' = \omega' \frac{z'^\perp}{|z'|} \quad \text{and} \quad \left(\frac{z'^\perp}{|z'|} \right)' = -\omega' \frac{z'}{|z'|}, \quad (4.13)$$

which hold for a curve z with sufficient regularity and its indicatrix ω . We take formally the derivative of the evolution equation in (4.3) and evaluate explicitly all the derivatives obtaining

$$\begin{aligned} \partial_t z'_\varepsilon + (r'_\varepsilon - (\lambda_\varepsilon \omega'_\varepsilon)') \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} - (r'_\varepsilon \omega'_\varepsilon - \lambda_\varepsilon \omega_\varepsilon'^2) \frac{z'_\varepsilon}{|z'_\varepsilon|} - \\ - (\lambda'_\varepsilon + (r'_\varepsilon \omega'_\varepsilon)') \frac{z'_\varepsilon}{|z'_\varepsilon|} - (\lambda''_\varepsilon \omega'_\varepsilon + r'_\varepsilon \omega_\varepsilon'^2) \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} = 0. \end{aligned} \quad (4.14)$$

We multiply by $z'_\varepsilon{}^\perp/|z'_\varepsilon|^2$ and $z'_\varepsilon/|z'_\varepsilon|$ and use $\partial_t \omega_\varepsilon = \partial_t z'_\varepsilon \cdot z'_\varepsilon{}^\perp/|z'_\varepsilon|^2$ and $\partial_t b_\varepsilon = \partial_t z'_\varepsilon \cdot z'_\varepsilon/|z'_\varepsilon|$ to obtain (1.13).

We present the proof of Theorem 1.2, which collects the regularity statements of Lemma 4.1, 4.2 and 4.3 and justifies the above computation in the distributional sense.

Proof. of Theorem 1.2. We start with a weak solution of (4.3) satisfying (4.5) and (4.8) respectively for all $v \in C_c^\infty(\mathbb{R}_+, C_s^\infty)$ according to Lemma 4.1. The boundary integrals imply that $\omega'_\varepsilon, b'_\varepsilon, \lambda_\varepsilon, r_\varepsilon|_{0,1} = 0$ a.e. on \mathbb{R}_+ .

We choose a regularising sequence η_k with $\text{supp} \eta_k \subset B(0, 1/k) \subset \mathbb{R}^2$ and set $v = \eta_k(\tilde{t} - t, \tilde{s} - s)$ for $(\tilde{t}, \tilde{s}) \in U_k$ with $U_k := [1/k, \infty) \times (1/k, 1 - 1/k)$ obtaining

$$\partial_t z_{\varepsilon,k} + \left(r_\varepsilon \frac{z_{\varepsilon,k}^\perp}{|z'_{\varepsilon,k}|} \right)' * \eta_k - \left(\lambda_\varepsilon \frac{z'_{\varepsilon,k}}{|z'_{\varepsilon,k}|} \right)' * \eta_k = 0,$$

where $z_{\varepsilon,k} := z_\varepsilon * \eta_k$. We omit the tilde and integrate against $-\left(\frac{z'_{\varepsilon,k}}{|z'_{\varepsilon,k}|} \varphi + \frac{z_{\varepsilon,k}^\perp}{|z'_{\varepsilon,k}|^2} \psi \right)'$ with $\varphi, \psi \in \mathcal{D}(\mathbb{R}_+ \times [0, 1])$ and k large enough such that U_k covers the support of both test functions,

$$\begin{aligned} & \iint_{U_k} \left[-\partial_t z_{\varepsilon,k} \cdot \left(z_{\varepsilon,k}^\perp \frac{1}{|z'_{\varepsilon,k}|^2} \psi \right)' - \right. \\ & \quad \left. - \left(\left(r_\varepsilon \frac{z_{\varepsilon,k}^\perp}{|z'_{\varepsilon,k}|} \right)' * \eta_k - \left(\lambda_\varepsilon \frac{z'_{\varepsilon,k}}{|z'_{\varepsilon,k}|} \right)' * \eta_k \right) \cdot \left(z_{\varepsilon,k}^\perp \frac{1}{|z'_{\varepsilon,k}|^2} \psi \right)' \right] ds dt = 0, \\ & \iint_{U_k} -\partial_t z_{\varepsilon,k} \cdot \left(\frac{z'_{\varepsilon,k}}{|z'_{\varepsilon,k}|} \varphi \right)' - \left(\left(r_\varepsilon \frac{z_{\varepsilon,k}^\perp}{|z'_{\varepsilon,k}|} \right)' * \eta_k - \left(\lambda_\varepsilon \frac{z'_{\varepsilon,k}}{|z'_{\varepsilon,k}|} \right)' * \eta_k \right) \cdot \left(\frac{z'_{\varepsilon,k}}{|z'_{\varepsilon,k}|} \varphi \right)' ds dt = 0. \end{aligned} \tag{4.15}$$

Making use of the regularity results in Theorem 1.1 it holds that $z'_{\varepsilon,k} \rightarrow z'_\varepsilon$ and, as a consequence, also $\omega_{\varepsilon,k} \rightarrow \omega_\varepsilon$, uniformly on compact subsets of $\mathbb{R}_+ \times [0, 1]$, where $\omega_{\varepsilon,k}$ is the indicatrix of $z_{\varepsilon,k}$ satisfying $z'_{\varepsilon,k}/|z'_{\varepsilon,k}| = (\cos(\omega_{\varepsilon,k}), \sin(\omega_{\varepsilon,k}))$. Using these results we perform integrations by parts with respect to s and t with the expressions in (4.15) that involve $\partial_t z_{\varepsilon,k}$ and pass to the limit, obtaining

$$\begin{aligned} \iint_{U_k} -\partial_t z_{\varepsilon,k} \cdot \left(\frac{z'_\varepsilon}{|z'_\varepsilon|} \varphi \right)' ds dt &= \iint_{U_k} \partial_t z'_{\varepsilon,k} \cdot \frac{z'_\varepsilon}{|z'_\varepsilon|} \varphi ds dt = \\ &= \iint_{U_k} -|z'_{\varepsilon,k}| \partial_t \varphi ds dt \rightarrow \iint_{U_k} -b_\varepsilon \partial_t \varphi ds dt \quad \text{as } k \rightarrow \infty \end{aligned}$$

and

$$\begin{aligned} \iint_{U_k} -\partial_t z_{\varepsilon,k} \cdot \left(z_{\varepsilon,k}^\perp \frac{1}{|z'_{\varepsilon,k}|^2} \psi \right)' ds dt &= \iint_{U_k} \partial_t z'_{\varepsilon,k} \cdot z_{\varepsilon,k}^\perp \frac{1}{|z'_{\varepsilon,k}|^2} \psi ds dt = \\ &= \iint_{U_k} \partial_t \omega_{\varepsilon,k} \psi ds dt = \iint_{U_k} -\omega_{\varepsilon,k} \partial_t \psi ds dt \rightarrow \\ &\rightarrow \int_0^\infty \int_0^1 -\omega_\varepsilon \partial_t \psi ds dt \quad \text{as } k \rightarrow \infty, \end{aligned}$$

where we used that $\partial_t \omega_{\varepsilon,k} = \partial_t z'_{\varepsilon,k} \cdot z_{\varepsilon,k}^\perp / |z'_{\varepsilon,k}|^2$. Making use of the fact that $z''_\varepsilon \in L_t^\infty L_s^2$ by Theorem 1.1 and $r'_\varepsilon, \lambda'_\varepsilon$ in $L_t^2 L_s^2$ by Lemma 4.2 we now pass to the limit

$k \rightarrow \infty$ in the remaining expressions of (4.15) and conclude

$$\int_0^\infty \int_0^1 \left[-\omega_\varepsilon \partial_t \psi - \left(r'_\varepsilon \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} - \lambda'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} - r_\varepsilon \omega'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} - \lambda_\varepsilon \omega'_\varepsilon \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} \right) \cdot \left(-\frac{z'_\varepsilon}{|z'_\varepsilon|} \omega'_\varepsilon \frac{\psi}{|z'_\varepsilon|} + \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} \left(\frac{\psi}{|z'_\varepsilon|} \right)' \right) \right] ds dt = 0,$$

$$\int_0^\infty \int_0^1 \left[-b_\varepsilon \partial_t \varphi - \left(r'_\varepsilon \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} - \lambda'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} - r_\varepsilon \omega'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} - \lambda_\varepsilon \omega'_\varepsilon \frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} \right) \cdot \left(\frac{z'_\varepsilon{}^\perp}{|z'_\varepsilon|} \omega'_\varepsilon \varphi + \frac{z'_\varepsilon}{|z'_\varepsilon|} \varphi' \right) \right] ds dt = 0.$$

This implies (1.20), (1.21) where every term is well defined by Lemma 4.2 and where we allow for ψ, ϕ in function spaces in which test functions are densely contained. \square

5. Energy dissipation Observe that the formal time derivative of the total energy as formulated in (1.10) is given by

$$\frac{d}{dt} \mathcal{E}_\varepsilon = \int_0^1 (b^2 \omega'_\varepsilon \partial_t \omega'_\varepsilon + \lambda_\varepsilon \partial_t b_\varepsilon) ds.$$

Now test the first equation in (1.13) formally against $(b_\varepsilon^2 \omega'_\varepsilon)' = b_\varepsilon r_\varepsilon$ obtaining

$$\int_0^1 b^2 \omega'_\varepsilon \partial_t \omega'_\varepsilon ds = \int_0^1 \left[-(r'_\varepsilon)^2 - r_\varepsilon^2 (\omega'_\varepsilon)^2 - \lambda'_\varepsilon \omega'_\varepsilon r_\varepsilon + \omega'_\varepsilon \lambda_\varepsilon r'_\varepsilon \right] ds$$

and the second one in (1.13) against λ_ε which yields

$$\int_0^1 \lambda_\varepsilon \partial_t b_\varepsilon ds = \int_0^1 \left[r'_\varepsilon \omega'_\varepsilon \lambda_\varepsilon - (r_\varepsilon \omega'_\varepsilon) \lambda'_\varepsilon - (\lambda'_\varepsilon)^2 - (\omega'_\varepsilon)^2 (\lambda_\varepsilon)^2 \right] ds.$$

Take the sum to conclude the energy dissipation equality (1.14) on a formal level.

This can be made rigorous stating that (1.14) holds weakly in time (Theorem 1.3).

Proof. of Theorem 1.3 Here the problem is that we cannot directly set ψ in (1.21) equal to $b_\varepsilon r'_\varepsilon$, since its time derivative cannot necessarily be interpreted as a function. Therefore we regularise with respect to t using a sequence of mollifiers $(\eta_k)_{k=1,2,\dots}$ with $\text{supp } \eta_k \subset [-1/k, 1/k]$.

For $\tilde{t} \geq 1/k$ we denote by $\omega_{\varepsilon,k}(\tilde{t}, s) := (\omega_\varepsilon *_{t} \eta_k)(\tilde{t}, s) = \int_0^\infty \omega_\varepsilon(t, s) \eta_k(\tilde{t} - t) dt$ the regularised version of ω_ε and evaluate (1.21) using $\psi(t, s) = \eta_k(\tilde{t} - t) (b_{\varepsilon,k}^2 \omega'_{\varepsilon,k})'(\tilde{t}, s)$. The time integral becomes part of the convolution and we find the following expression which holds pointwise for every $\tilde{t} \geq 1/k$,

$$\int_0^1 \left[\partial_t \omega_{\varepsilon,k} (b_{\varepsilon,k}^2 \omega'_{\varepsilon,k})' - \left((r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon) \frac{1}{b_\varepsilon} \right) *_{t} \eta_k (b_{\varepsilon,k} r_{\varepsilon,k})' + \left((r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon) \frac{b'_\varepsilon}{b_\varepsilon^2} \right) *_{t} \eta_k b_{\varepsilon,k} r_{\varepsilon,k} - \left((r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) \frac{\omega'_\varepsilon}{b_\varepsilon} \right) *_{t} \eta_k b_{\varepsilon,k} r_{\varepsilon,k} \right] ds = 0.$$

Observe that we write the convolution of various expressions and η_k explicitly using the symbol $*_t$. Furthermore, set $\phi = \eta_k(\tilde{t}-t)\lambda_{\varepsilon,k}(\tilde{t},s)$ in (1.20) where $\lambda_{\varepsilon,k} = b_{\varepsilon,k}(\omega'_{\varepsilon,k})^2 - b''_{\varepsilon,k} + \frac{1}{\varepsilon}E'(b_{\varepsilon,k})$,

$$\int_0^1 [\partial_t b_{\varepsilon,k} \lambda_{\varepsilon,k} + (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) *_t \eta_k \lambda'_{\varepsilon,k} + ((\omega'_\varepsilon \lambda_\varepsilon - r'_\varepsilon) \omega'_\varepsilon) *_t \eta_k \lambda_{\varepsilon,k}] ds = 0.$$

We omit the tildes, take the sum of the two equations above and integrate against the test function in time $\vartheta \in \mathcal{D}(\mathbb{R}_+)$ obtaining

$$\begin{aligned} 0 = \int_{1/k}^\infty & \left[-\partial_t \vartheta \mathcal{E}_{\varepsilon,k} + \vartheta \int_0^1 \left[\left((r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon) \frac{1}{b_\varepsilon} \right) *_t \eta_k (b_{\varepsilon,k} r_{\varepsilon,k})' + \right. \right. \\ & \left. \left. + \left(-(r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon) \frac{b'_\varepsilon}{b_\varepsilon^2} + (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) \frac{\omega'_\varepsilon}{b_\varepsilon} \right) *_t \eta_k b_{\varepsilon,k} r_{\varepsilon,k} + \right. \right. \\ & \left. \left. + (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) *_t \eta_k \lambda'_{\varepsilon,k} + ((\omega'_\varepsilon \lambda_\varepsilon - r'_\varepsilon) \omega'_\varepsilon) *_t \eta_k \lambda_{\varepsilon,k} \right] ds \right] dt \quad \forall \tilde{t} \geq 1/k, \quad (5.1) \end{aligned}$$

where $\mathcal{E}_{\varepsilon,k} = \int_0^1 \left[\frac{1}{2} (\omega'_{\varepsilon,k} b_{\varepsilon,k}^2 + (b'_{\varepsilon,k})^2) + \frac{1}{\varepsilon} E(b_{\varepsilon,k}) \right] ds$ and k large enough such that $\text{supp } \vartheta \subset [1/k, \infty)$. Next we use the fact that all the convolved terms like $\omega'_{\varepsilon,k}$, $r'_{\varepsilon,k}$, $(r_\varepsilon (\omega'_\varepsilon)^2) *_t \eta_k$, etc. converge strongly in $L^2_{t,\text{loc}} L^2_s$ to their original counterparts, which is illustrated in more detail in the proof of Theorem 4 in [Oel10]. Hence we pass to the limit as $k \rightarrow \infty$ in (5.1) obtaining

$$\begin{aligned} \int_0^\infty \partial_t \vartheta \mathcal{E}_\varepsilon dt &= \int_0^\infty \left[\vartheta \int_0^1 \left[\left((r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon) \frac{1}{b_\varepsilon} \right) (b_\varepsilon r'_\varepsilon)' + \right. \right. \\ & \left. \left. + \left(-(r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon) \frac{b'_\varepsilon}{b_\varepsilon^2} + (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) \frac{\omega'_\varepsilon}{b_\varepsilon} \right) b_\varepsilon r'_\varepsilon + \right. \right. \\ & \left. \left. + (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon) \lambda'_\varepsilon + ((\omega'_\varepsilon \lambda_\varepsilon - r'_\varepsilon) \omega'_\varepsilon) \lambda_\varepsilon \right] ds \right] dt \quad \forall \tilde{t} \geq 1/k = \\ &= \int_0^\infty \vartheta \int_0^1 \left[(r'_\varepsilon - \lambda_\varepsilon \omega'_\varepsilon)^2 + (r_\varepsilon \omega'_\varepsilon + \lambda'_\varepsilon)^2 \right] ds dt \quad \forall \tilde{t} \geq 0, \end{aligned}$$

which is a weak in time formulation of (1.14).

The second equivalent formulation in (1.15) is then a consequence of the following computation based on (4.1) and (4.13).

$$\begin{aligned} \left(z''''_\varepsilon - \left(\frac{1}{\varepsilon} E'(|z'_\varepsilon|) \frac{z'_\varepsilon}{|z'_\varepsilon|} \right)' \right)^2 &= \left(\left(r_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} + \lambda_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \right)' \right)^2 \\ &= \left(r'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} + \lambda'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} - r_\varepsilon \omega'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} + \lambda_\varepsilon \omega'_\varepsilon \frac{z'_\varepsilon}{|z'_\varepsilon|} \right)^2 \\ &= (r'_\varepsilon + \lambda_\varepsilon \omega'_\varepsilon)^2 + (\lambda'_\varepsilon - r_\varepsilon \omega'_\varepsilon)^2, \end{aligned}$$

which concludes the proof. \square

6. Long time convergence One of the main tools in this section will be the fact that the best constant in the Poincaré-type inequality

$$\int_0^1 v^2 ds \leq C \int_0^1 v'^2 ds \quad (6.1)$$

for $v \in H_0^1((0,1))$ is given by the reciprocal value of the first eigenvalue of the differential operator v'' in that space, $C = 1/\pi^2$.

As a first step in proving coercivity of \mathcal{D}_ε with respect to \mathcal{E}_ε we find

LEMMA 6.1. *Let \mathcal{E}_ε be the total energy defined in (1.1) and (1.10) respectively and let r_ε and λ_ε as defined in (1.12), then there is a constant $C_1 > 0$ such that*

$$\mathcal{E}_\varepsilon \leq C_1 (\|r_\varepsilon\|_{L^2_s}^2 + \|\lambda_\varepsilon\|_{L^2_s}^2).$$

Proof. We will treat the three different expressions in (1.10) separately. First observe that

$$\int_0^1 (b_\varepsilon \omega'_\varepsilon)^2 d\tilde{s} \leq \frac{1}{\pi^2} \int_0^1 ((b_\varepsilon \omega'_\varepsilon)')^2 d\tilde{s} \leq \frac{b_{\max}^2}{\pi^2} \int_0^1 r_\varepsilon^2 d\tilde{s},$$

which immediately implies that

$$\|\omega'_\varepsilon\|_{L^2}^2 \leq \frac{b_{\max}^2}{\pi^2 b_{\min}^2} \int_0^1 r_\varepsilon^2 d\tilde{s}. \quad (6.2)$$

Next we use the definition of λ_ε in (1.12) which we multiply by $(b_\varepsilon - 1)$ obtaining $-(b_\varepsilon - 1)b'_\varepsilon \leq (b_\varepsilon - 1)\lambda_\varepsilon + b_\varepsilon(\omega'_\varepsilon)^2$ since $(b-1)\frac{1}{\varepsilon}E'(b) \geq 0$ as E is convex and takes its minimum at $x=1$. We integrate and using $b'_\varepsilon(0) = b'_\varepsilon(1) = 0$ we obtain

$$\begin{aligned} \int_0^1 (b'_\varepsilon)^2 d\tilde{s} &\leq \int_0^1 [(b_\varepsilon - 1)\lambda_\varepsilon + b_\varepsilon(\omega'_\varepsilon)^2] d\tilde{s} \leq \\ &\leq \max(b_{\max} - 1, 1 - b_{\min}) \|\lambda_\varepsilon\|_{L^2}^2 + b_{\max} \|\omega'_\varepsilon\|_{L^2}^2. \end{aligned} \quad (6.3)$$

Finally, to obtain a control on $\int_0^1 \frac{1}{\varepsilon} E(b_\varepsilon) d\tilde{s}$ we use Lemma 2.5 and (4.12), which implies that

$$\frac{1}{\varepsilon} \|E'(b_\varepsilon)\|_{L^2}^2 \leq \varepsilon \|\lambda_\varepsilon - b_\varepsilon(\omega'_\varepsilon)^2\|_{L^2}^2.$$

The fact that $\omega''_\varepsilon = \frac{r_\varepsilon - 2b'_\varepsilon \omega'_\varepsilon}{b_\varepsilon}$ allows to estimate

$$\begin{aligned} \|\omega''_\varepsilon\|_{L^\infty}^2 &= \left\| \int_0^s \omega''_\varepsilon d\tilde{s} \right\|_{L^\infty}^2 \leq \|\omega''_\varepsilon\|_{L^1}^2 \leq \\ &\leq \frac{2}{b_{\min}^2} (\|r_\varepsilon\|_{L^1}^2 + 4\|b'_\varepsilon\|_{L^2}^2 \|\omega'_\varepsilon\|_{L^2}^2) \leq \frac{2}{b_{\min}^2} (\|r_\varepsilon\|_{L^1}^2 + 8b_{\min}^{-2} \mathcal{E}_I \|b'_\varepsilon\|_{L^2}^2), \end{aligned} \quad (6.4)$$

where we used (3.1). Summarising, now using Lemma 2.5, we obtain

$$\begin{aligned} \int_0^1 \frac{1}{\varepsilon} E(b_\varepsilon) d\tilde{s} &\leq \frac{1}{2\kappa} \int_0^1 \frac{1}{\varepsilon} E'(b_\varepsilon)^2 d\tilde{s} \leq \frac{1}{2\kappa} \varepsilon \|\lambda_\varepsilon - b_\varepsilon(\omega'_\varepsilon)^2\|_{L^2}^2 \leq \\ &\leq \varepsilon \frac{1}{\kappa} (\|\lambda_\varepsilon\|_{L^2}^2 + b_{\max}^2 \|(\omega'_\varepsilon)^2\|_{L^2}^2) \leq \varepsilon \frac{1}{\kappa} (\|\lambda_\varepsilon\|_{L^2}^2 + b_{\max}^2 \|\omega'_\varepsilon\|_{L^\infty}^2 \|\omega'_\varepsilon\|_{L^2}^2) \leq \\ &\leq \varepsilon \frac{1}{\kappa} (\|\lambda_\varepsilon\|_{L^2}^2 + 2b_{\min}^{-2} \mathcal{E}_I b_{\max}^2 \|\omega'_\varepsilon\|_{L^\infty}^2). \end{aligned} \quad (6.5)$$

Now apply the inequalities (6.2), (6.3) and (6.5) combined with (6.4) to the total energy as formulated in (1.10) to obtain the result. \square

We finally conclude the exponential decay of the energy formulated in Theorem 1.4.

Proof. of Theorem 1.4 We start with the result of Lemma 6.1 and obtain

$$\begin{aligned}
\mathcal{E}_\varepsilon &\leq C_1 \int_0^1 (r_\varepsilon^2 + \lambda_\varepsilon^2) ds = \\
&= C_1 \int_0^1 \left[\left(z_\varepsilon''' \cdot \frac{z_\varepsilon'^\perp}{|z_\varepsilon'|} \right)^2 + \left(z_\varepsilon''' \cdot \frac{z_\varepsilon'}{|z_\varepsilon'|} - \frac{1}{\varepsilon} E(|z_\varepsilon'|) \right)^2 \right] ds = \\
&= C_1 \int_0^1 \left| z_\varepsilon''' - \frac{1}{\varepsilon} E(|z_\varepsilon'|) \frac{z_\varepsilon'}{|z_\varepsilon'|} \right|^2 ds \leq \\
&\leq \frac{C_1}{\pi^2} \int_0^1 \left| \left(z_\varepsilon''' - \frac{1}{\varepsilon} E(|z_\varepsilon'|) \frac{z_\varepsilon'}{|z_\varepsilon'|} \right)' \right|^2 ds = \frac{C_1}{\pi^2} \mathcal{D}_\varepsilon,
\end{aligned} \tag{6.6}$$

where we used (1.15). As a consequence, the energy dissipation \mathcal{D}_ε is coercive with respect to the energy and the following Poincaré-type inequality holds

$$\mathcal{D}_\varepsilon / \mathcal{E}_\varepsilon \geq \pi^2 / C_1, \tag{6.7}$$

which implies directly the result. \square

As an immediate consequence of Theorem 1.4 we obtain the following refined bounds.

COROLLARY 6.2. *Using the definitions in (1.16) we have*

$$\|\partial_t z_\varepsilon\|_{L_t^2 L_s^2}^2 = \|m_\varepsilon\|_{L_t^2 L_s^2}^2 + \|n_\varepsilon\|_{L_t^2 L_s^2}^2 = \mathcal{E}_I \tag{6.8}$$

As a consequence it holds that

$$\|m_\varepsilon\|_{L_t^2 L_s^2} \leq \sqrt{\mathcal{E}_I}, \quad \|n_\varepsilon\|_{L_t^2 L_s^2} \leq \sqrt{\mathcal{E}_I}.$$

We also obtain sharper bounds on the following quantities,

$$\begin{aligned}
&\|r_\varepsilon\|_{L_t^2 L_s^\infty}, \|\lambda_\varepsilon\|_{L_t^2 L_s^\infty} \leq \sqrt{\mathcal{E}_I}, \\
&\|\omega'_\varepsilon \lambda_\varepsilon\|_{L_t^2 L_s^2}, \|\omega'_\varepsilon r_\varepsilon\|_{L_t^2 L_s^2} \leq \sqrt{2\mathcal{E}_I} b_{\min}^{-1} \quad \text{and} \\
&\|\lambda'_\varepsilon\|_{L_t^2 L_s^2}, \|r'_\varepsilon\|_{L_t^2 L_s^2} \leq \sqrt{2\mathcal{E}_I} b_{\min}^{-1} + \sqrt{\mathcal{E}_I}.
\end{aligned}$$

Proof. Integrate (1.14) with respect to time and combine it with (4.10). Using this result we go again through the estimates in the proof of Lemma 4.2. \square

7. Convergence as $\varepsilon \rightarrow 0$

LEMMA 7.1. *Let $(\omega_\varepsilon, b_\varepsilon, \lambda_\varepsilon)$ be a solution to (1.13) according to Theorem 1.2, then there is ω_0 such that $\omega_\varepsilon \rightarrow \omega_0$ in $L^1((0, T), W^{2,r}(0, 1)) \cap L^q((0, T), H^1(0, 1))$, $1 \leq r, q < \infty$.*

Proof. We obtain the strong convergence of ω_ε by an application of the Lions-Aubin Lemma as formulated in [Sim87]. On the one hand, using (1.17), observe that

the family of distributions $\{\partial_t \omega_\varepsilon\}$ is uniformly bounded in $L^1(0, T; H^{-1}((0, 1)))$ as

$$\begin{aligned} \langle \partial_t \omega_\varepsilon, v \rangle_{H_s^{-1}} &= \int_0^1 \left[n_\varepsilon \left(\frac{v'}{b_\varepsilon} - \frac{v}{b_\varepsilon^2} b'_\varepsilon \right) + m_\varepsilon \omega'_\varepsilon \frac{v}{b_\varepsilon} \right] ds \leq \\ &\leq \frac{1}{b_{\min}} \left(\int_0^1 |n_\varepsilon| |v'| ds + \frac{\|v\|_{L_s^\infty}}{b_{\min}} \int_0^1 |n_\varepsilon| |b'_\varepsilon| ds + \|v\|_{L_s^\infty} \int_0^1 |\omega'_\varepsilon| |m_\varepsilon| ds \right) \end{aligned}$$

for all $v \in H_0^1((0, 1))$. This implies

$$\begin{aligned} \int_0^T \left| \sup_{v \in H_0^1} \frac{\langle \partial_t \omega_\varepsilon, v \rangle_{H_s^{-1}}}{\|v\|_{H_0^1}} \right| dt &\leq \sup_{v \in H_0^1} \frac{1}{b_{\min} \|v\|_{H_0^1}} \left(\int_0^T \int_0^1 |n_\varepsilon| |v'| ds dt + \right. \\ &\quad \left. + \frac{\|v\|_{L_s^\infty}}{b_{\min}} \int_0^T \int_0^1 |n_\varepsilon| |b'_\varepsilon| ds dt + \|v\|_{L_s^\infty} \int_0^T \int_0^1 |\omega'_\varepsilon| |m_\varepsilon| ds dt \right) \leq \\ &\leq \frac{1}{b_{\min}} \left(\sqrt{T} \|n_\varepsilon\|_{L^2(0, T; L_s^2)} + \frac{1}{b_{\min}} \|b'_\varepsilon\|_{L^2(0, T; L_s^2)} \|n_\varepsilon\|_{L^2(0, T; L_s^2)} + \right. \\ &\quad \left. + \|\omega'_\varepsilon\|_{L^2(0, T; L_s^2)} \|m_\varepsilon\|_{L^2(0, T; L_s^2)} \right) \leq \\ &\leq \frac{\sqrt{T}}{b_{\min}} \left(\|n_\varepsilon\|_{L^2(0, T; L_s^2)} + \sqrt{2} \mathcal{E}_I \left(\frac{1}{b_{\min}} \|n_\varepsilon\|_{L^2(0, T; L_s^2)} + b_{\min}^{-2} \|m_\varepsilon\|_{L^2(0, T; L_s^2)} \right) \right). \end{aligned}$$

Here we used the uniform bounds of (3.1) and Corollary 6.2.

Since the family $\{\omega_\varepsilon\}$ is uniformly bounded in $L^1(0, T; W^{3,1}(0, 1))$ by Lemma 4.3 and due to the compact embedding $W^{1,1}((0, 1)) \hookrightarrow L^r((0, 1))$ ($1 \leq r < \infty$) it holds that the family ω_ε is relatively compact in $L^1(0, T; W^{2,r}(0, 1))$.

By interpolation, using the uniform bound in $L_t^\infty H^1(0, 1)$ on ω_ε , this holds also true in $L^q(0, T; H^1(0, 1))$ for any $1 \leq q < \infty$. \square

LEMMA 7.2. *Let $(\omega_\varepsilon, b_\varepsilon, \lambda_\varepsilon)$ be a solution to (1.13) according to Theorem 1.2, then it holds that $b_\varepsilon \rightarrow 1$ in $L^1((0, T), W^{1,r}(0, 1)) \cap L^q((0, T), H^1(0, 1))$, $1 \leq r, q < \infty$.*

Proof. We obtain the strong convergence of b_ε by an application of the Lions-Aubin Lemma as formulated in [Sim87]. On the one hand, using (1.13) and the definitions (1.16), observe that the family of distributions $\{\partial_t b_\varepsilon\}$ is uniformly bounded in $L^1(0, T; H^{-1}((0, 1)))$ as

$$\langle \partial_t b_\varepsilon, v \rangle_{H_s^{-1}} = \int_0^1 [-m_\varepsilon v' + n_\varepsilon \omega'_\varepsilon v] ds \leq \int_0^1 |m_\varepsilon| |v'| ds + \|v\|_{L_s^\infty} \int_0^1 |\omega'_\varepsilon| |n_\varepsilon| ds$$

for all $v \in H_0^1((0, 1))$. This implies

$$\begin{aligned} \int_0^T \left| \sup_{v \in H_0^1} \frac{\langle \partial_t b_\varepsilon, v \rangle_{H_s^{-1}}}{\|v\|_{H_0^1}} \right| dt &\leq \\ &\leq \sup_{v \in H_0^1} \frac{1}{\|v\|_{H_0^1}} \left(\int_0^T \int_0^1 |m_\varepsilon| |v'| ds dt + \|v\|_{L_s^\infty} \int_0^T \int_0^1 |\omega'_\varepsilon| |n_\varepsilon| ds dt \right) \leq \\ &\leq \int_0^T \|m_\varepsilon\|_{L_s^2} dt + \|\omega'_\varepsilon\|_{L^2(0, T; L_s^2)} \|n_\varepsilon\|_{L^2(0, T; L_s^2)} \leq \\ &\leq \sqrt{T} \|n_\varepsilon\|_{L^2(0, T; L_s^2)} + \sqrt{T 2 b_{\min}^{-2} \mathcal{E}_I} \|m_\varepsilon\|_{L^2(0, T; L_s^2)}. \end{aligned}$$

Here we used the uniform bounds of (3.1) and Corollary 6.2.

Since the family $\{b_\varepsilon\}$ is uniformly bounded in $L^1(0, T; W^{2,1}(0, 1))$ by Lemma 4.3 and due to the compact embedding $W^{1,1}((0, 1)) \hookrightarrow L^r((0, 1))$ ($1 \leq r < \infty$) it holds that the family b_ε is relatively compact in $L^1(0, T; W^{1,r}(0, 1))$.

By interpolation, using the uniform bound in $L_t^\infty H^1(0, 1)$ on b_ε , this holds also true in $L^q(0, T; H^1(0, 1))$ for any $1 \leq q < \infty$.

Finally, we consider the first inequality formulated in Lemma 2.5 and obtain as a consequence of energy dissipation (1.14) that

$$\|b_\varepsilon(t, \cdot) - \mathbf{1}\|_{L_s^2} \leq \frac{2}{\kappa} \sqrt{\varepsilon \mathcal{E}_I} \quad \text{for all } t > 0, \quad (7.1)$$

which allows to identify the limit of any strongly converging subsequence of the family $\{b_\varepsilon\}$ as the constant function $\mathbf{1}$.

□

LEMMA 7.3. *Let $(\omega_\varepsilon, b_\varepsilon, \lambda_\varepsilon)$ be a solution to (1.13) according to Theorem 1.2, then there are subsequences and limit functions such that $\lambda_\varepsilon \rightharpoonup \lambda_0$ in $L_t^2 H_s^1$ and $r_\varepsilon \rightarrow \omega_0''$ in $L^2((0, T), L_s^2)$ with $r_\varepsilon \rightarrow \omega_0''$ in $L_t^2 H_s^1$.*

Proof. The strong convergence of ω_ε and b_ε (Lemma 7.1 and 7.2) imply the strong convergence of r_ε to ω_0'' . The weak convergence of r_ε' and also the weak convergence of λ_ε to a limit function λ_0 are a consequence of the uniform bounds stated in Corollary 6.2. □

Finally we finish with the proof of Theorem 1.5 which summarises the results of Lemma 7.1, 7.2 and 7.3 and states the consistency of the model (1.5) with the limit model (1.6) and, as a consequence, the strong convergence of m_ε and n_ε .

Proof. of Theorem 1.5 The convergence results of Lemma 7.1, 7.2 and 7.3 immediately allow to pass to the limit as $\varepsilon \rightarrow 0$ in the system (1.20), (1.21) obtaining as the limit system (2.2), (2.3).

Observe that the definitions (1.16) together with the strong convergence of ω_ε (Lemma 7.1) and the weak convergence of r_ε and λ_ε (Lemma 7.3) imply that

$$m_\varepsilon \rightarrow \omega_0' \omega_0'' + \lambda_0' \quad \text{and} \quad n_\varepsilon \rightarrow \omega_0''' - \omega_0' \lambda_0 \quad \text{in} \quad L_t^2 L_s^2.$$

The fact that the equality (6.8) also holds in the limit as $\varepsilon_j \rightarrow 0$, (2.7), implies

$$\|m_\varepsilon\|_{L_t^2 L_s^2}^2 + \|n_\varepsilon\|_{L_t^2 L_s^2}^2 = \mathcal{E}_I = \|\omega_0' \omega_0'' + \lambda_0'\|_{L_t^2 L_s^2}^2 + \|\omega_0''' - \omega_0' \lambda_0\|_{L_t^2 L_s^2}^2.$$

These two facts imply the strong convergence of m_ε and n_ε since

$$\begin{aligned} & \int_0^\infty \int_0^1 [(m_\varepsilon - m_0)^2 + (n_\varepsilon - n_0)^2] ds dt = \\ & = \int_0^\infty \int_0^1 [m_\varepsilon^2 - 2m_\varepsilon m_0 + m_0^2 + n_\varepsilon^2 - 2n_\varepsilon n_0 + n_0^2] ds dt \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

□

Acknowledgement.

This work has been supported by the Vienna Science and Technology Fund (WWTF) through the project "Mathematical modeling of actin driven cell migration" of C. Schmeiser.

REFERENCES

- [AGS05] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré, *Gradient flows in metric spaces and in the space of probability measures*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2005. MR MR2129498
- [BCM10] C. Bost, G.-H. Cottet, and E. Maitre, *Convergence analysis of a penalization method for the three-dimensional motion of a rigid body in an incompressible viscous fluid*, SIAM J. Numer. Anal. **48** (2010), no. 4, 1313–1337. MR 2684337
- [CMR09] Eduardo Casas, Mariano Mateos, and Jean-Pierre Raymond, *Penalization of Dirichlet optimal control problems*, ESAIM Control Optim. Calc. Var. **15** (2009), no. 4, 782–809. MR 2567245 (2010h:49063)
- [DGMT80] Ennio De Giorgi, Antonio Marino, and Mario Tosques, *Problems of evolution in metric spaces and maximal decreasing curve*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) **68** (1980), no. 3, 180–187. MR MR636814 (83m:49052)
- [Dil92] Ellis Harold Dill, *Kirchhoff's theory of rods.*, Arch. Hist. Exact Sci. **44** (1992), no. 1, 1–23 (English).
- [HY03] X.X. Huang and X.Q. Yang, *Convergence analysis of a class of nonlinear penalization methods for constrained optimization via first-order necessary optimality conditions*, Journal of Optimization Theory and Applications **116** (2003), 311–332, 10.1023/A:1022503820909.
- [Kir59] G. R. Kirchhoff, *Über das Gleichgewicht und die Bewegung eines unendlich dünnen elastischen Stabes.*, J. Reine Angew. Math. **56** (1859), 285–313.
- [Lin89] Anders Linnér, *Some properties of the curve straightening flow in the plane*, Trans. Amer. Math. Soc. **314** (1989), no. 2, 605–618. MR MR989580 (90g:58022)
- [Oel10] Dietmar Oelz, *On the curve straightening flow of inextensible, open, planar curves*, SēMA Journal (2010), no. 54, 5–24.
- [OS10] Dietmar Oelz and Christian Schmeiser, *Cell mechanics: from single scale-based models to multiscale modelling*, ch. How do cells move? mathematical modelling of cytoskeleton dynamics and cell migration, Chapman and Hall / CRC Press, 2010.
- [OS11] ———, *Simulation of lamellipodia with arbitrary shape.*, J. Math. Biol., Epub ahead of print. (2011).
- [Sim87] Jacques Simon, *Compact sets in the space $L^p(0,T;B)$* , Ann. Mat. Pura Appl. (4) **146** (1987), 65–96. MR MR916688 (89c:46055)