

Fermionic Solution of the Andrews–Baxter–Forrester Model. II. Proof of Melzer’s Polynomial Identities¹

S. Ole Warnaar²

Received August 29, 1995

We compute the one-dimensional configuration sums of the ABF model using the fermionic technique introduced in part I of this paper. Combined with the results of Andrews, Baxter, and Forrester, we prove polynomial identities for finitizations of the Virasoro characters $\chi_{h, a}^{(r, \sigma)}(q)$ as conjectured by Melzer. In the thermodynamic limit these identities reproduce Rogers–Ramanujan-type identities for the unitary minimal Virasoro characters conjectured by the Stony Brook group. We also present a list of additional Virasoro character identities which follow from our proof of Melzer’s identities and application of Bailey’s lemma.

KEY WORDS: ABF model; one-dimensional configuration sums; Fermi lattice gas; Melzer’s polynomial identities; Rogers–Ramanujan identities; Virasoro characters.

1. INTRODUCTION

Probably among the most celebrated results in mathematics are the identities of Rogers and Ramanujan,^(1–3)

$$\sum_{m=0}^{\infty} \frac{q^{m(m+a)}}{(q)_m} = \frac{1}{(q)_{\infty}} \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j+1+2a)/2}, \quad a=0, 1 \quad (1.1)$$

where $(q)_m = \prod_{k=1}^m (1 - q^k)$, $m > 0$, and $(q)_0 = 1$. In the context of modern physics, one recognizes the right-hand side of these identities to be the Rocha-Caridi expression for the Virasoro characters $\chi_{1, 2-a}^{(2, 5)}(q)$ of minimal conformal field theory $M(2, 5)$.⁽⁴⁾ As such, the Rogers–Ramanujan identities can be seen as character identities of some Virasoro algebra. A natural

¹ Dedicated to the memory of Piet Kasteleyn.

² Mathematics Department, University of Melbourne, Parkville, Victoria 3052, Australia; e-mail: warnaar@maths.mu.oz.au.

question is whether the other Virasoro characters also admit identities of the Rogers–Ramanujan type. For the important class of *unitary* minimal models $M(r - 1, r)$, this was answered affirmative in a remarkable paper by the Stony Brook group.^{(5), 2} However, the results of ref. 5 were all based on extensive numerical studies, and actual proofs remained elusive.

Among the many methods of proof of the original Rogers–Ramanujan identities an elegant approach is that of first proving the polynomial identities^(8, 9)

$$\sum_{m=0}^{\infty} q^{m(m+a)} \begin{bmatrix} L-m-a \\ m \end{bmatrix} = \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j+1+2a)/2} \begin{bmatrix} L \\ \lfloor \frac{1}{2}(L-5j-a) \rfloor \end{bmatrix} \tag{1.2}$$

for all $L \geq a$. Here $\lfloor x \rfloor$ denotes the integer part of x and $\begin{bmatrix} N \\ m \end{bmatrix}$ is a Gaussian polynomial defined as

$$\begin{bmatrix} N \\ m \end{bmatrix} = \begin{cases} (q)_N / (q)_m (q)_{N-m} & 0 \leq m \leq N \\ 0 & \text{otherwise} \end{cases} \tag{1.3}$$

Clearly, in the limit of $L \rightarrow \infty$ we recover the Rogers–Ramanujan identity (1.1). To prove the *finitized* Rogers–Ramanujan identities (1.2) it suffices to check that both left- and right-hand sides satisfy the elementary recurrences $f_L = f_{L-1} + q^{L-1} f_{L-2}$ as well as the same initial conditions for $L = a, a + 1$.

In an attempt to find proofs of the identities for the characters $\chi_{b,a}^{(r-1,r)}(q)$ (see next section for their actual form), Melzer followed Schur’s approach and conjectured finitizations similar to those in (1.2). However, Melzer’s polynomial identities were sufficiently complicated not to lead to a straightforward proof using recurrences. It was only after Melzer proved the cases $r = 3$ (Ising) and $r = 4$ (tricritical Ising)⁽¹⁰⁾ that Berkovich succeeded in proving recurrences for the polynomial identities for all $\chi_{b,1}^{(r,r-1)}(q)$.⁽¹¹⁾

In this paper we present a combinatorial proof for Melzer’s identities, based on yet another observation made by Melzer. Again the motivation for this has been the original Rogers–Ramanujan identities (1.1), whose finitization (1.2) can be viewed as evaluations of the sum

$$\sum_{\substack{\sigma_1, \dots, \sigma_{L-1} = 0, 1 \\ \sigma_j \sigma_{j+1} = 0}} q^{\sum_{k=1}^{L-1} k \sigma_k}, \quad \sigma_0 = a, \quad \sigma_L = 0 \tag{1.4}$$

² By now character identities of Rogers–Ramanujan type for all minimal Virasoro characters $\chi^{(p,p')}(q)$ have been found.⁽⁵⁻⁷⁾

in two intrinsically different ways. Similar to this, Melzer has argued that the polynomial identities for the finitized $\chi_{b,a}^{(r-1,r)}(q)$ characters arise from computing the sums

$$X_L(a, b) = \sum_{\substack{\sigma_1, \dots, \sigma_{L-1} = 0 \\ |\sigma_{j+1} - \sigma_j| = 1}}^{r-2} q^{\sum_{k=1}^L k|\sigma_{k+1} - \sigma_{k-1}|/4},$$

$$\sigma_0 = a - 1, \quad \sigma_L = b - 1, \quad \sigma_{L+1} = b \quad (1.5)$$

for all $a = 1, \dots, r - 1$ and $b = 1, \dots, r - 2$.

We will take this observation as the starting point for proving the polynomial and Rogers–Ramanujan identities for the (finitized) characters $\chi_{b,a}^{(r-1,r)}(q)$. That is, we give two different methods to compute (1.5), one leading to a so-called *fermionic* expression similar to the left-hand side of (1.2) and one method leading to a so-called *bosonic* expression similar to the right-hand side of (1.2). In fact, it should be noted that $X_L(a, b)$ defined above is exactly the *one-dimensional configuration sum* $X_L(a, b, c)$, with $c = b + 1$, as defined by Andrews, Baxter, and Forrester in their computation of the order parameters of the $(r - 1)$ -state ABF model in regime III.⁽¹²⁾ Hence, computing the sum (1.5) amounts to computing the order parameters of the ABF model. The fact that (finitized) Rogers–Ramanujan identities arise from calculating order parameters of solvable lattice models is in fact not new, and indeed the sum (1.4) is exactly the one encountered by Baxter in his solution of the hard-hexagon model in regime I.⁽¹³⁾

The remainder of this paper is organized as follows. In the next section we describe Melzer’s polynomial identities, their limiting Rogers–Ramanujan-type form, and some other Virasoro character identities that follow from the proof of Melzer’s identities and the application of the Andrews–Bailey construction.^(14–16) Then, in Section 3 we compute the configuration sums (1.5) using the technique developed in part I of this paper.⁽¹⁷⁾ This amounts to reinterpreting the sum (1.5) as the grand canonical partition function of a one-dimensional gas of charged particles obeying certain Fermi-type exclusion rules. In Section 4 we describe the original approach of ABF for computing (1.5) using recurrence relations. Together with the result of Section 3, this proves Melzer’s polynomial identities. We finally end with a discussion of our result and an outlook on related problems and generalizations.

To end this introduction we make some further remarks on the problem described in this paper. First, as mentioned before, an altogether different kind of proof of Melzer’s identities has recently been given for the case of $\chi_{b,1}^{(r-1,r)}(q)$ by Berkovich.⁽¹¹⁾ This method of proof, which in fact is

applicable to all unitary minimal characters,⁽⁷⁾ is based on recursive instead of combinatorial arguments.³

Second, in their solution of the ABF model, Andrews, Baxter, and Forrester also considered the configuration sums $X_L(a, b, c)$, with $c = b - 1$. Hence, to compute complete all configuration sums of the ABF model, more general sums than those defined in (1.5) have to be considered. However, from simple symmetry arguments^(12, 10) (see also Section 3) one can easily deduce that computing (1.5) suffices to obtain expressions for all $X_L(a, b, c)$.

Finally we remark that Melzer⁽¹⁰⁾ and Kedem *et al.*⁽⁵⁾ conjecture (in the general case) four fermionic expressions for each (finitized) character. In this paper we give detailed proof of only two of the four. For the remaining two representations we did not succeed in finding a derivation in terms of a Fermi lattice-gas.

2. MELZER'S POLYNOMIAL IDENTITIES AND RELATED ROGERS–RAMANUJAN IDENTITIES

In this section we give a summary of identities proven by the calculations carried out in Sections 3 and 4. First we describe the polynomial identities conjectured by Melzer⁽¹⁰⁾ and their limiting Rogers–Ramanujan-type form as discovered by the Stony Brook group.⁽⁵⁾ Then we list two classes of character identities for nonunitary minimal models which, as recently pointed out by Foda and Quano,⁽⁶⁾ arise from Melzer's identities and the Andrews–Bailey construction.^(14–16)

2.1. Identities for the (Finitized) Virasoro Characters

$$\chi_{b, a}^{(r-1, r)}(q)$$

Before we state the polynomial identities as conjectured by Melzer, we need some notation. We denote the incidence matrix of the A_{r-3} Dynkin diagram by \mathcal{I} , with $\mathcal{I}_{j, k} = \delta_{j, k-1} + \delta_{j, k+1}$, $j, k = 1, \dots, r-3$. The Cartan matrix of A_{r-3} is denoted as C , and is related to \mathcal{I} by $C_{j, k} = 2\delta_{j, k} - \mathcal{I}_{j, k}$. We also define the $(r-3)$ -dimensional (column) vectors \mathbf{m} and \mathbf{e}_j , $j = 1, \dots, r-3$, by $(\mathbf{m})_j = m_j$ and $(\mathbf{e}_j)_k = \delta_{j, k}$, and set $m_0 = m_{r-2} = 0$, $\mathbf{e}_0 = \mathbf{e}_{r-2} = \mathbf{0}$. With this notation, using the Gaussian polynomials as defined in (1.3), we can state Melzer's conjectures as the following identities for $a = 1, \dots, r-1$, $b = 1, \dots, r-2$, and $L - |a - b| \in 2\mathbb{Z}_{\geq 0}$,⁴

³ Berkovich has subsequently proven Melzer's identities for all characters, but his results remain unpublished.⁽¹⁸⁾

⁴ Throughout this paper we use the notation $x \equiv y$ to mean $x \equiv y \pmod{2}$. Also, the sums $\sum_{x \equiv y}$ and \sum_x are shorthand notations for $\prod_j \sum_{x_j \geq 0; x_j \equiv y_j}$ and $\prod_j \sum_{x_j \geq 0}$, respectively.

$$\begin{aligned}
 f_{a,b} & \sum_{\mathbf{m} \equiv \mathbf{Q}_{a,b}} q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \prod_{j=1}^{r-3} \left[\frac{1}{2}(\mathcal{J} \mathbf{m} + L \mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-b-1})_j \right] \\
 & = \sum_{j=-\infty}^{\infty} \left\{ q^{j(r-1)j+rb-(r-1)a} \left[\begin{array}{c} L \\ \frac{1}{2}(L+a-b)-rj \end{array} \right] \right. \\
 & \quad \left. - q^{((r-1)j+b)(rj+a)} \left[\begin{array}{c} L \\ \frac{1}{2}(L-a-b)-rj \end{array} \right] \right\} \quad (2.1)
 \end{aligned}$$

with $f_{a,b} = q^{-(a-b)(a-b-1)/4}$ and

$$\mathbf{Q}_{a,b} = \mathbf{Q}_{a,b}^{(r-3)} = (\mathbf{e}_{r-a-2} + \mathbf{e}_{r-a-4} + \cdots) + (\mathbf{e}_{r-b-2} + \mathbf{e}_{r-b-4} + \cdots) \quad (2.2)$$

We note that in our derivation of the left-hand side of (2.1) in Section 3, this restriction naturally arises in the following form, (mod 2)-equivalent to (2.2):

$$(\mathbf{Q}_{a,b})_j = \min(a-1, r-j-2) + \min(b-1, r-j-2) \quad (2.3)$$

In ref. 10, yet another expression for the left-hand side of (2.1) was conjectured as

$$f_{a,b} \sum_{\mathbf{m} \equiv \mathbf{R}_{a,b}} q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{a-1}} \prod_{j=1}^{r-3} \left[\frac{1}{2}(\mathcal{J} \mathbf{m} + L \mathbf{e}_1 + \mathbf{e}_{a-1} + \mathbf{e}_{r-b-1})_j \right] \quad (2.4)$$

where

$$\mathbf{R}_{a,b} = (r-a-1) \boldsymbol{\rho} + (\mathbf{e}_a + \mathbf{e}_{a+2} + \cdots) + (\mathbf{e}_{r-b-2} + \mathbf{e}_{r-b-4} + \cdots) \quad (2.5)$$

with $\boldsymbol{\rho} = \sum_{j=1}^{r-3} \mathbf{e}_j$. Clearly, for $a=1$ and for $a=r-1$ the fermionic expressions in (2.1) and (2.4) coincide.

As mentioned in the introduction, we have no explanation of this alternative fermionic form in terms of a Fermi gas, and (2.4) is listed only for completeness.

Taking the finitization parameter L to infinity, we find that (2.1) leads to Rogers–Ramanujan-type identities for unitary minimal Virasoro characters. Here we recall the well-known Rocha-Caridi expression for all (normalized) characters $\chi_{r,s}^{(p,p')}(q)$ of minimal CFT $M(p,p')$,

$$\chi_{r,s}^{(p,p')}(q) = \frac{1}{(q)_{\infty}} \sum_{j=-\infty}^{\infty} \{ q^{j(pp'j+p'r-ps)} - q^{(jp+r)(jp'+s)} \} \quad (2.6)$$

for $r = 1, \dots, p - 1, s = 1, \dots, p' - 1$, with p and p' coprime. We thus find that the right-hand side of (2.1) gives the *bosonic* Rocha-Caridi expression for $\chi_{b,a}^{(r-1,r)}(q)$, whereas the left-hand side leads to a *fermionic* counterpart,

$$\chi_{b,a}^{(r-1,r)}(q) = f_{a,b} \sum_{\mathbf{m} \in \mathbf{Q}_{a,b}} \frac{q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}}}{(q)_{m_1}} \times \prod_{j=2}^{r-3} \left[\frac{\frac{1}{2}(\mathcal{J} \mathbf{m} + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-b-1})_j}{m_j} \right] \tag{2.7}$$

This result is one of the many celebrated conjectures for fermionic character representations made by the Stony Brook group; see, e.g., refs. 5, 19, and 20.

An obvious symmetry of (2.6) is $\chi_{r,s}^{(p,p')}(q) = \chi_{p-r,p'-s}^{(p,p')}(q)$. Making the transformation $a \rightarrow r - a$ and $b \rightarrow r - b - 1$ in the fermionic expression (2.7), we see that this symmetry is not at all manifest, except for $b = 1$ and $a = 1, r - 2$. Hence we have two different fermionic representations for each character of the unitary minimal series.

To end our discussion on Melzer’s polynomial identities, we remark that in ref. 10 identities were also given for finitizations of the characters $\chi_{b,a}^{(r-1,r)}(q)$, with finitization parameter L such that $L + a - b \neq 0$. Since these can simply be obtained from (2.1) and (2.4) by the above-mentioned symmetry transformation, they are not listed here as separate identities.

2.2. Rogers–Ramanujan Identities for $\chi_{a,(k+1)b}^{(r,(k+1)r-1)}(q)$ and $\chi_{b,(k+1)a}^{(r-1,(k+1)r-k)}(q)$

It was recently pointed out by Foda and Quano⁽⁶⁾ that many new Virasoro character identities can be obtained by applying some powerful lemmas proven by Bailey and Andrews to Melzer’s polynomial identities. The main idea of these lemmas is to prove the more complicated Rogers–Ramanujan-type identities by showing that they are a consequence of identities that are easier to prove. Here we will not state the relevant lemmas but refer the interested reader to the work of Foda and Quano⁽⁶⁾ and to the original work of Bailey^(14,15) and Andrews.⁽¹⁶⁾

In both series of Virasoro character identities given below, we encounter the k by k matrix B with entries $B_{j,l} = \min(j, l)$. We note that this matrix is the inverse of the Cartan-type matrix of the tadpole graph with k nodes; $(B^{-1})_{j,l} = 2\delta_{j,l} - \mathcal{J}_{j,l}^{(k)}$, with incidence matrix of the tadpole graph given by $\mathcal{J}_{j,l}^{(k)} = \delta_{j,l-1} + \delta_{j,l+1} + \delta_{j,l}\delta_{j,k}$, $j, l = 1, \dots, k$. We will also use the k -dimensional vectors \mathbf{n} and $\boldsymbol{\epsilon}_k$, whose j th entries read n_j and $\delta_{k,j}$, respectively.

2.2.1. $\chi_{a, (k+1)b}^{(r, (k+1)r-1)}(\mathbf{q})$. Substituting the *Bailey pair* read off from (2.1) into the *Bailey chain* of length k , we obtain

$$\begin{aligned} &\chi_{a, (k+1)b}^{(r, (k+1)r-1)}(\mathbf{q}) \\ &\stackrel{(a \equiv b)}{=} f_{a, b} q^{-k(a-b)^2/4} \sum_{\mathbf{n}} \sum_{\mathbf{m} \equiv \mathbf{Q}_{a, b}} \frac{q^{\mathbf{n}^T B \mathbf{n}}}{(q)_{n_1} \cdots (q)_{n_{k-1}} (q)_{2n_k}} \\ &\quad \times q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \prod_{j=1}^{r-3} \left[\frac{\frac{1}{2}(\mathcal{J} \mathbf{m} + 2n_k \mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-b-1})_j}{m_j} \right] \\ &\stackrel{(a \not\equiv b)}{=} f_{a, b} q^{-k((a-b)^2-1)/4} \sum_{\mathbf{n}} \sum_{\mathbf{m} \equiv \mathbf{Q}_{a, b}} \frac{q^{\mathbf{n}^T B(\mathbf{n} + \boldsymbol{\varepsilon}_k)}}{(q)_{n_1} \cdots (q)_{n_{k-1}} (q)_{2n_k + 1}} \\ &\quad \times q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \prod_{j=1}^{r-3} \left(\frac{\frac{1}{2}(\mathcal{J} \mathbf{m} + (2n_k + 1)\mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-b-1})_j}{m_j} \right) \end{aligned} \tag{2.8}$$

valid for all $k \geq 1$, $a = 1, \dots, r-1$, $b = 1, \dots, r-2$.

The proof of this result for $a=1$ was first noted by Foda and Quano,⁽⁶⁾ using the proof of Melzer’s identities for $a=1$ as established by Berkovich.⁽¹¹⁾ The fermionic expression in (2.8) can also be found in ref. 7.

2.2.2. $\chi_{b, (k+1)a}^{(r-1, (k+1)r-k)}(\mathbf{q})$. Substitute the *dual Bailey pair* obtained from (2.1) into the *Bailey chain* of length $k+1$. Then make the change of variables $m_j \rightarrow m_{j+1}$, followed by $2n_{k+1} + |a-b| \rightarrow m_1$, $n_k \rightarrow n_k + \frac{1}{2}(m_1 - |a-b|)$, and $r \rightarrow r-1$. Finally, interchanging a and b , then using

$$(\mathbf{Q}_{a, b}^{(r-3)})_j \equiv \begin{cases} (\mathbf{Q}_{b, a}^{(r-4)})_{j-1}, & j = 2, \dots, r-3 \\ a-b, & j = 1 \end{cases} \tag{2.9}$$

true for $a = 1, \dots, r-3$, $b = 1, \dots, r-2$, yields

$$\begin{aligned} &\chi_{b, (k+1)a}^{(r-1, (k+1)r-k)}(\mathbf{q}) \\ &= f_{a, b} q^{-k(a-b)^2/4} \sum_{\mathbf{n}} \sum_{\mathbf{m} \equiv \mathbf{Q}_{a, b}} \frac{q^{(\mathbf{n} + (1/2)m_1 \boldsymbol{\varepsilon}_k)^T B(\mathbf{n} + (1/2)m_1 \boldsymbol{\varepsilon}_k)}}{(q)_{n_1} \cdots (q)_{n_k}} \\ &\quad \times \frac{q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}}}{(q)_{m_1}} \prod_{j=2}^{r-3} \left[\frac{\frac{1}{2}(\mathcal{J} \mathbf{m} + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-b-1})_j}{m_j} \right] \end{aligned} \tag{2.10}$$

valid for all $k \geq 0$, $a = 1, \dots, r-3$, $b = 1, \dots, r-2$. Note that for $k=0$, corresponding to a *Bailey chain* of length 1, we actually recover a subset of the character identities (2.7) for $M(r-1, r)$.

For $a = b = 1$, (2.10) was conjectured in ref. 5. The proof for $a = 1$ can again be found in ref. 6, though the actual form of the fermionic side therein rather differs due to the sequence of the transformations carried out above. The fermionic form (2.10) can also be found in ref. 7.

3. FERMIONIC SOLUTION OF THE ABF MODEL

We now come to the main part of this paper, the evaluation of the one-dimensional configuration sums (1.5) of the ABF model. This yields, up to the prefactor $f_{a,b}$, the left-hand side of the identity (2.1). To establish this, we first reformulate the sum (1.5) as the generating function of certain restricted lattice paths. We then compute this generating function by identifying each path as a configuration of charged fermions on a one-dimensional lattice. This identification allows us to view $X_L(a,b)$ as the grand-canonical partition function of a one-dimensional Fermi gas. Because of the one-dimensional nature of this gas, its partition function can readily be computed.

3.1. Restricted Lattice Paths

To reformulate the sum (1.5) in terms of lattice paths, we first give some basic definitions.

Definition 1. An ordered sequence of spins $\{\sigma_0, \sigma_1, \dots, \sigma_{L+1}\}$ is called *admissible* if:

- $\sigma_j \in \{0, 1, \dots, r-2\}$ for $j=0, \dots, L+1$
- $|\sigma_{j+1} - \sigma_j| = 1$ for $j=0, \dots, L$, and
- $\sigma_0 = a-1$, $\sigma_L = b-1$, and $\sigma_{L+1} = b$

Definition 2. Let $\{\sigma_0, \sigma_1, \dots, \sigma_{L+1}\}$ be an admissible sequence of spins. Plot all pairs (j, σ_j) in the (x, y) -plane and interpolate between each pair of neighboring points by a straight line segment. The resulting graph is called a *restricted lattice path*.

An example of a restricted lattice path for $a = 3$ and $b = 5$ is shown in Fig. 1.

To write the one-dimensional configuration sum as a sum over restricted lattice paths, first notice that the restrictions on the σ 's in (1.5) precisely correspond to those defining an admissible sequence of spins. Consequently, each restricted lattice path corresponds to one of the terms in the sum (1.5) and, conversely, each term in the sum corresponds to a




Fig. 1. An example of a restricted lattice path in $\text{rlp}(0, r)$.

restricted lattice path. Given an admissible sequence, its total weight is decomposed as follows. If $\sigma_{j-1} < \sigma_j < \sigma_{j+1}$ or $\sigma_{j-1} > \sigma_j > \sigma_{j+1}$ this contributes a factor $q^{j/2}$ and if $\sigma_{j-1} < \sigma_j > \sigma_{j+1}$ or $\sigma_{j-1} > \sigma_j < \sigma_{j+1}$ this contributes a factor 1. In terms of the restricted lattice paths this simply means that for each integer point j along the x axis we get a factor 1 if (j, σ_j) is an extremum and a factor $q^{j/2}$ otherwise. Here the terminals of a path are to be viewed as extrema. Writing this in the language of statistical mechanics, we get, setting $q = \exp(-\beta)$,

$$X_L(a, b) = \sum_{\text{restricted lattice paths}} \exp \left[-\beta \sum_{j=1}^L E(j) \right] \tag{3.1}$$

with energy function E given by

$$E(j) = \begin{cases} 0 & \text{if the path has an extremum at } (x \text{ position}) j \\ \frac{1}{2}j & \text{otherwise} \end{cases} \tag{3.2}$$

Each of the lattice paths in the sum (3.1) starts in $(0, a-1)$, ends in $(L, b-1)$, $(L+1, b)$, and is restricted to the strip $0 \leq y \leq r-2$. We now define $\text{rlp}(\mu, r)$ as the set of all restricted lattice paths with minimal y value equal to μ and maximal y value less than or equal to $r-2$. Hence we can write

$$X_L(a, b) = \sum_{\mu=0}^{\min(a, b) - 1} \Xi_L(a, b; \mu, r) \tag{3.3}$$

with

$$\Xi_L(a, b; \mu, r) = \sum_{\text{rlp}(\mu, r)} \exp \left[-\beta \sum_{j=1}^L E(j) \right] \tag{3.4}$$

Noting the obvious relation $\mathcal{E}_L(a, b; \mu, r) = \mathcal{E}_L(a - \mu, b - \mu; 0, r - \mu)$ gives

$$X_L(a, b) = \sum_{\mu=0}^{\min(a, b)-1} \mathcal{E}_L(a - \mu, b - \mu; 0, r - \mu) \quad (3.5)$$

and we conclude that to compute $X_L(a, b)$ it suffices to compute the sum (3.4) for $\mu=0$ and arbitrary a, b , and r .

So far we only have reformulated the problem of computing $X_L(a, b)$, and it is by no means clear that $\mathcal{E}_L(a, b) := \mathcal{E}_L(a, b; 0, r)$ is any simpler to evaluate than (1.5). To make some real progress, we will show in the next section that $\mathcal{E}_L(a, b)$ can be viewed as the grand canonical partition function of a one-dimensional gas of charged fermions. In other words, each path in $\text{rlp}(0, r)$ can be viewed as a configuration of an appropriately defined Fermi gas. Now decomposing the sum over all Fermi-gas configurations into a sum over the configuration with fixed particle content (FC) and a sum over the particle content (C), we get

$$\mathcal{E}_L(a, b) = \sum_C Z(C; a, b) \quad (3.6)$$

with $Z(C; a, b)$ the partition function of the one-dimensional Fermi gas,

$$Z(C; a, b) = \sum_{\text{FC}} \exp \left[-\beta \sum_{j=1}^L E(j) \right] \quad (3.7)$$

3.2. A One-Dimensional Fermi Gas

To interpret each restricted lattice path in $\text{rlp}(0, r)$ as a configuration of particles, we need some more terminology. In fact, since some of the concepts introduced below are somewhat awkward to describe, but easily explained pictorially, we state some definitions purely graphically.

In the previous section restricted lattice paths were introduced as paths from $(0, a-1)$ to $(L, b-1)$, $(L+1, b)$, restricted to the strip $0 \leq y \leq r-2$, such that $y_{j+1} - y_j = \pm 1$ for all consecutive points (j, y_j) and $(j+1, y_{j+1})$ on the path. We somewhat relax these conditions by defining a *lattice path* as follows.

Definition 3. A *lattice path* is a restricted lattice path with arbitrary (integer) begin and endpoint.

In particular, if a lattice path ends in (j, y_j) , the y -coordinate of the second-last point can either be $y_j - 1$ or $y_j + 1$.




Fig. 2. Typical examples of left-boundary, bulk, and right-boundary complexes.

We use the previous definition to define a very important object, a *complex*.⁵ This will be used subsequently to decompose each restricted lattice path into particles.

Definition 4. A *bulk complex* is a lattice path from (j, y_j) to (k, y_k) , with (j, y_j) and (k, y_k) connected by a dashed horizontal line, such that $y_j = y_k, y_l > y_j$ for all $j < l < k$.

A *left-boundary complex* is a lattice path from $(0, a - 1)$ to $(j, 0)$, such that $y_k > 0$ for all $k < j$, and with $(0, 0)$ and $(j, 0)$ connected by a horizontal dashed line and $(0, 0)$ and $(0, a - 1)$ connected by a vertical solid line.

A *right-boundary complex* is a lattice path from $(j, 0)$ to $(L + 1, b - 1)$, $(L + 1, b)$, such that $y_k > 0$ for all $k > j$ and with $(L + 1, 0)$ and $(j, 0)$ connected by a horizontal dashed line and $(L + 1, 0)$ and $(L + 1, b)$ connected by a vertical solid line.

Examples of left-boundary, bulk, and right-boundary complexes can be found in Fig. 2. With respect to the above definition, we remark that the term *complex* is chosen since we wish to view each complex as a collection of charged particles moved on top of each other. To make this explicit, we define particles in the following two definitions.

Definition 5. A *pure bulk particle* of charge j is a bulk complex with a single local maximum of height j (measured with respect to its dashed line).

A *pure left-boundary particle* of charge $(a - 1)/2$ is a left-boundary complex with a single local maximum, located at $(0, a - 1)$.

A *pure right-boundary particle* of charge $b/2$ is a right-boundary complex with a single local maximum.

⁵ In ref. 21, Bressoud gives a lattice path interpretation of the Andrews–Gordon generalizations of the Roger–Ramanujan identities.^(22, 23) In Bressoud’s terminology a complex corresponds to a *mountain*.

The graphical representation of pure particles is given in Fig. 3.

To introduce the more general idea of a particle, we need some simple terminology.

- The *peak* of a bulk complex is the leftmost highest point. Similarly, the peak of a particle is its highest point.
- The *origin* of a particle or complex is the left- and downmost point; the *endpoint* of a particle or complex is the right- and downmost point; the *baseline* of a particle or complex is the dashed line connecting the begin and endpoint.
- The *contour* of a particle or complex is its part drawn with solid lines.

Using this terminology, we present the following definitions.

Definition 6. A *bulk particle* of charge j is a pure bulk particle of charge j whose contour is interrupted at arbitrary integer points by horizontal dashed lines of even length.

A *left boundary particle* of charge $(a-1)/2$ is a pure left-boundary particle of charge $(a-1)/2$ whose contour to the right of $(0, a-1)$ is interrupted at arbitrary integer points by horizontal dashed lines of even length.

A *right-boundary particle* of charge $b/2$ is a pure right-boundary particle of charge $b/2$ whose contour to the left of $(L, b-1)$ is interrupted at arbitrary integer points by horizontal dashed lines of even length.

Typical examples of particles are shown in Fig. 4. We note that for later convenience the contour of the boundary particles is drawn with thicker lines than that of the bulk particles.

With the above set of definitions we now give a prescription to divide each restricted lattice path into particles. This will be done by giving an algorithm that divides a complex into a particle and several smaller complexes. Each of these new complexes is either a particle or is again divided into a particle and yet smaller complexes. This procedure is continued until




Fig. 3. The graphical representation of pure particles. The charges are, from left to right, $(a-1)/2$, j , and $b/2$, respectively.




Fig. 4. Typical examples of left-boundary, bulk, and right-boundary particles. The charges of the particles are $(a-1)/2$, j , and $b/2$, respectively.

the entire complex is divided into particles. Since each lattice path can trivially be divided into complexes, this gives a procedure to divide any restricted lattice path into particles.

(0) Draw a dashed line along the x axis from $(0, 0)$ to $(L+1, 0)$ and draw bold lines from $(0, 0)$ to $(0, a-1)$ and $(L+1, 0)$ to $(L+1, b)$. This divides each restricted lattice path into a left-boundary complex, a right-boundary complex, and a number of bulk complexes. For the restricted lattice path of Fig. 1, we get, for example, four complexes, two of which are of bulk type. If $a = 1$, the left-boundary complex is absent.

Now consider each of the complexes obtained above. If such a complex is a particle (in which case it is pure), we are done with it. If not, go to step (1) in case of a bulk complex and to (1_L) and (1_R) in case of left- and right-boundary complexes, respectively.

(1) Start at the peak of the complex and move down to the right along the contour till the endpoint of the complex. When a local minimum is reached, i.e., the contour starts going up again, we draw a dashed line from this local minimum to the right until we cross the contour. At that point we move further down along the contour. If another minimum occurs we repeat the above, etc.

Repeat the above, now moving to the left. That is, start from the peak of the complex and move down to the left till the origin of the complex. If a local minimum is reached, we draw a dashed line to the left and continue our movement down when the dashed line intersects the contour.

As a result of the above step we have divided the complex into a particle (which is not pure) and several (at least one) smaller complexes. The peak and the baseline of the particle are the peak and the baseline of the original complex. Now go to (2).

(1_L) Start from $(0, a-1)$. Move to the right of this point down along the contour of the complex till its endpoint. If a local minimum is reached [which could be the point $(0, a-1)$ itself], draw a dashed line from this

steps. Letting n_j denote the number of bulk particles of charge j , we thus have the completeness relation

$$a + b - 1 + 2 \sum_{j=1}^{r-2} j n_j = L + 1 \quad (3.8)$$

Using this relation, we can compute n_{r-2} given the occupation numbers n_1, \dots, n_{r-3} . For this reason (and anticipating things to come), we define the column vector $\mathbf{n} = {}^T(n_1, \dots, n_{r-3})$, and when we say “a restricted lattice path has particle content $C = \mathbf{n}$,” we mean by this the particle content $C = \{n_1, \dots, n_{r-2}\}$ subject to the restriction (3.8).

Having associated a configuration of particles with each path in $\text{rlp}(0, r)$, we define $\text{rlp}(\mathbf{n})$ as the subset of paths in $\text{rlp}(0, r)$ with particle content \mathbf{n} . This puts us in a position to properly define what we mean by the Fermi-gas partition function as introduced in (3.6),

$$Z(C; a, b) = Z(\mathbf{n}; a, b) = \sum_{\text{rlp}(\mathbf{n})} \exp \left[-\beta \sum_{j=1}^L E(j) \right] \quad (3.9)$$

with energy function defined in (3.2).

So far, we have repeatedly used the term Fermi gas, without any clear motivation. Clearly, we have defined all allowed configurations of our one-dimensional system of charged particles, as well its Hamiltonian or energy function, but the actual nature of the system remains rather elusive. However, in our actual computation of Z in the next subsection it turns out to be expedient to define rules of motion that allow one to obtain any configuration with content \mathbf{n} from a given so-called minimal configuration with the same content. These rules of motion have a clear fermionic character, in that particles of the same charge cannot exchange position, unlike particles of different charge.

3.3. Computation of $Z(\mathbf{n}; a, b)$

In this section we compute the partition function of the one-dimensional Fermi gas. Throughout the section we assume the particle content to be \mathbf{n} .

To compute the sum over all particle configurations, we first select a particular configuration called the *minimal configuration*.⁷ It will be defined purely graphically.

⁷ From a statistical mechanics point of view *ground-state configuration* may be more appropriate, but we prefer to conform to our earlier naming in ref. 17.

Using the above three results, we compute the energy of the minimal configuration as

$$\begin{aligned}
 E_{\min} &= E_a + E_b + \sum_{j=1}^{r-2} \sum_{l=1}^{n_j} E_j \left(a - 2 + \min(b, r - j - 1) \right. \\
 &\quad \left. + 2j(l - 1) + 2 \sum_{k=j+1}^{r-2} kn_k \right) \\
 &= \sum_{j=1}^{r-2} (j - 1) n_j \left(jn_j + 2 \sum_{k=j+1}^{r-2} kn_k \right) + (a + b - 2) \sum_{j=1}^{r-2} (j - 1) n_j \\
 &\quad + \sum_{j=r-b}^{r-2} (b - r + j + 1) n_j + \frac{1}{4}(a - 1)(a - 2) \\
 &\quad + \frac{1}{4}(b - 1)(2a + b - 2)
 \end{aligned} \tag{3.13}$$

To simplify this expression, we eliminate n_{r-3} using the completeness relation (3.8). This yields

$$\begin{aligned}
 E_{\min} &= \sum_{j=1}^{r-3} \left(\sum_{k=l}^j \frac{k(r-j-2)}{r-2} + \sum_{k=j+1}^{r-3} \frac{j(r-k-2)}{r-2} \right) n_j n_k \\
 &\quad - \left(\sum_{j=1}^{r-b-1} \frac{j(b-1)}{r-2} + \sum_{j=r-b}^{r-3} \frac{(r-b-1)(r-j-2)}{r-2} + L \sum_{j=1}^{r-3} \frac{r-j-2}{r-2} \right) n_j \\
 &\quad + \frac{L^2(r-3) + 2L(b-1) - (a-1)(r-a-1) + (b-1)(r-b-1)}{4(r-2)}
 \end{aligned} \tag{3.14}$$

We now recall the definition of the inverse Cartan matrix of the Lie algebra A_{r-3} ,

$$C_{j,k}^{-1} = \begin{cases} k(r-j-2)/(r-2), & k \leq j \\ j(r-k-2)/(r-2) & k \geq j \end{cases} \tag{3.15}$$

Using this, we finally obtain the following result.

Lemma 1. The energy of the minimal configuration is given by

$$\begin{aligned}
 E_{\min} &= \sum_{j,k=1}^{r-3} \left(n_j - \frac{L}{2} \delta_{j,1} - \frac{1}{2} \delta_{j,r-b-1} - \frac{1}{2} \delta_{j,r-a-1} \right) C_{j,k}^{-1} \\
 &\quad \times \left(n_k - \frac{L}{2} \delta_{k,1} - \frac{1}{2} \delta_{k,r-b-1} + \frac{1}{2} \delta_{k,r-a-1} \right)
 \end{aligned} \tag{3.16}$$

3.3.2. Contribution of the Nonminimal Configurations. To compute the contribution to the partition function of the other configurations, we define rules of motion which generate all nonminimal configurations from the minimal one. These rules break up into several different *elementary moves* as follows.

Definition 8. Let $X = \{(x_1, y_1), (x_2, y_2), (x_3, y_3), (x_4, y_4)\}$ denote a sequence of four points on the contour of a configuration, each pair of consecutive points connected straight lines, such that the contour in between (x_1, y_1) and (x_4, y_4) does not belong to a boundary particle. We may then replace this sequence by a new sequence of four points as follows:

Move L_u : If $y_4 \leq y_2 < y_3 < y_1$,

$$L_u(X) = \{(x_1, y_1), (x_2 - 1, y_2 + 1), (x_3 - 1, y_3 + 1), (x_4, y_4)\}$$

Move R_d : If $y_4 < y_2 < y_3 \leq y_1$,

$$R_d(X) = \{(x_1, y_1), (x_2 + 1, y_2 - 1), (x_3 + 1, y_3 - 1), (x_4, y_4)\}$$

Move L_d : If $y_1 < y_3 < y_2 \leq y_4$,

$$L_d(X) = \{(x_1, y_1), (x_2 - 1, y_2 - 1), (x_3 - 1, y_3 - 1), (x_4, y_4)\}$$

Move R_u : If $y_1 \leq y_3 < y_2 < y_4$,

$$R_u(X) = \{(x_1, y_1), (x_2 + 1, y_2 + 1), (x_3 + 1, y_3 + 1), (x_4, y_4)\}$$

Besides these “bulk-type” moves we need some special boundary moves.

Definition 9. Let $X = \{(x_1, y_1), (x_2, y_2), (x_3, y_3), (x_4, y_4)\}$ be four points on the contour of a configuration, each pair of consecutive points connected by a straight line. We may then replace X as follows:

Move L'_u : Let $(x_1, y_1) = (x_2 - 1, y_2 + 1)$. If $y_2 = y_4 < y_3 < r - 2$ and the contour between the first two points belongs to the left-boundary particle,

$$L'_u(X) = \{(x_2 - 1, y_2 + 1), (x_3 - 1, y_3 + 1), (x_4 - 1, y_4 + 1), (x_4, y_4)\}$$

where the contour between the last two points belongs to the left-boundary particle.

Move R'_d : Let $(x_4, y_4) = (x_3 + 1, y_3 - 1)$. If $y_1 = y_3 < y_2$ and the contour between the first two points belongs to the left-boundary particle,

$$R'_d(X) = \{(x_1, y_1), (x_1 + 1, y_1 - 1), (x_2 + 1, y_2 - 1), (x_3 + 1, y_3 - 1)\}$$

where the contour between the last two points belongs to the left-boundary particle.

Move L'_d : Let $(x_1, y_1) = (x_2 - 1, y_2 - 1)$. If $y_2 = y_4 < y_3$ and the contour between the first two points belongs to the right-boundary particle,

$$L'_d(X) \{(x_2 - 1, y_2 - 1), (x_3 - 1, y_3 - 1), (x_4 - 1, y_4 - 1), (x_4, y_4)\}$$

where the contour between the last two points belongs to the right-boundary particle.

Move R'_u : Let $(x_4, y_4) = (x_3 + 1, y_3 + 1)$. If $y_1 = y_3 < y_2 < r - 2$, $y_3 < b - 1$, and the contour between the last two points belongs to the right-boundary particle,

$$R'_u(X) = \{(x_1, y_1), (x_1 + 1, y_1 + 1), (x_2 + 1, y_2 + 1), (x_3 + 1, y_3 + 1)\}$$

where the contour between the first two points belongs to the right-boundary particle.

For the graphical interpretation of this long list of moves, see Fig. 7.

To fully appreciate these moves, we list its main characteristics in several lemmas, which are at the core of our fermionic computation of the one-dimensional configuration sums.

Lemma 2. The elementary moves are reversible. That is, if there is a move of type M_s^p from a configuration C to a configuration C' , then there is a move of type \bar{M}_s^p from C' to C . Here $M = L$ or R ; $s = u$ or d ; $p = \cdot, ' \text{ or } ''$; and $\bar{R} = L, \bar{L} = R, \bar{u} = d$, and $\bar{d} = u$.





Fig. 7. (a) The moves L_u and R_d . (b) The moves L_d and R_u . (c) The moves L'_u and R'_d . (d) The moves L'_d and R'_u .

Proof. Let us show this for L_u . The other moves follow in similar manner. Let X be a sequence of four extrema as in Definition 8, satisfying $y_4 \leq y_2 < y_3 < y_1$. Hence we can carry out L_u to obtain $X' = L_u(X) = \{(x'_1, y'_1), (x'_2, y'_2), (x'_3, y'_3), (x'_4, y'_4)\}$. From the definition of the move L_u , we find that $y'_4 \leq y'_2 - 1 < y'_3 - 1 < y_1$. We rewrite this to obtain $y'_4 < y'_2 < y'_3 < y'_1$ and hence we can carry out the move R_d to obtain $R_d(X') = X$. ■

Lemma 3. The moves leave the particle content \mathbf{n} fixed.

Proof. This follows immediately from the graphical representation of the moves shown in Fig. 7, where the dashed lines represent the baselines of the pure particles being moved. Note here that the graphical representations of the moves R_u and L_d are the generic cases. Performing a move of type R_u to a sequence X as defined in Definition 8, with $y_2 = y_4 - 1$, may lead to a “jump” of the baseline. A similar thing may happen when performing a move of type L_d to a sequence with $y_2 = y_4$:



Lemma 4. Given the minimal configuration, we cannot make any of the R -type moves.

Proof. we can only make moves of type R_d if we have a sequence of X as in Definition 8, with $y_4 < y_2$. Clearly this does not occur. We can only make moves of type R_u if we have a sequence X , with $y_2 < y_4$. Again this does not occur. We cannot make a move of type R'_d , since the left-boundary particle is in its pure form. Finally, we cannot make a move of type R'_u , since all particles of charge $j \geq r - b - 1$ have their peak at $y = r - 2$, and all particles of charge $j \leq r - b - 1$ have their endpoint at $y = b - 1$. ■

Lemma 5. If a configuration is not the minimal one, we can always make a move of type R .

Proof. By construction the minimal configuration is the only configuration that does not meet any of the conditions required for one of the R -type moves. In particular, all maxima (apart from the initial point of the path) are of decreasing order and all minima of increasing order. This completely fixes the path. If one of these two properties is broken somewhere along the path, we can always make an R -type move. ■

These first four lemmas can be combined to give the following proposition:

Proposition 1. All nonminimal paths are generated by moves of type L from the minimal configuration. All nonminimal configurations can be reduced to the minimal configuration by moves of type R .

Having established the above proposition, we can perform the actual calculation of the generation function \mathcal{G} of the moves of type L . Again we prepare some lemmas to obtain the desired result.

Lemma 6. Each move of type L generates a factor q .

Proof. We show this for the typical case of move L_u . The total energy E of a sequence of extrema X is

$$E = \sum_{\substack{j=x_1+1 \\ j \neq x_2, x_3}}^{x_4-1} j/2 \tag{3.17}$$

Similarly, the energy E' of the sequence $X' = L_u(X)$ is

$$E' = \sum_{\substack{j=x_1+1 \\ j \neq x_2-1, x_3-1}}^{x_4-1} j/2 \tag{3.17}$$

Hence we find

$$e^{-\beta(E' - E)} = q^{E' - E} = q \quad \blacksquare \tag{3.19}$$

In the following it will be convenient to label the bulk particles in the minimal configuration, letting $p_{j,l}$ denote the l th particle of charge j , counted from the left. To now generate all nonminimal configurations, we give an ordering for carrying out the moves of type L .

- The particle $p_{j,l}$ is moved to the left using moves of type L , prior to any of the particles $p_{k,m}$, with $k \leq j$, and with $m > l$ if $k = j$.

Assuming this order (which will be justified later), we have the following result.

Lemma 7. The maximal number of L -type moves $p_{j,1}$ can make is

$$m_j = 2 \sum_{k=j+1}^{r-2} (k-j) n_k + \min(a-1, r-j-2) + \min(b-1, r-j-2) \tag{3.20}$$

Proof. We prove this lemma in two steps. In the first step (3.20) is shown to be true for the minimal configuration, and in the second step it is shown that m_j is invariant under having moved the particles $p_{k,m}$, with $k > j$, prior to $p_{j,1}$.

Let us start to calculate the number of L -type moves needed to exchange the position of two particles of charge k and j , $k > j$, with j positioned immediately to the right of k . In such a configuration of two particles we have a sequence $X = \{(x_1, y_1), \dots, (x_5, y_5)\}$ of points connected by straight lines, with $y_1 = y_3 = y_5$ and $y_2 = k$ and $y_4 = j$. From these conditions it follows that move L_u can be carried out $k - j$ times to the sequence $\{(x_2, y_2), \dots, (x_5, y_5)\}$. This gives a new sequence $X' = \{(x'_1, y'_1), \dots, (x'_5, y'_5)\}$, with $y'_1 = y'_5, y'_2 = y'_4 = k$, and $y'_3 = k - j$. From these conditions it follows that move L_d can be carried out $k - j$ times to the sequence $\{(x'_1, y'_1), \dots, (x'_4, y'_4)\}$. This gives the final sequence $X'' = \{(x''_1, y''_1), \dots, (x''_5, y''_5)\}$, with $y''_1 = y''_3 = y''_5, y''_2 = j$, and $y''_4 = k$. The total number of moves carried out is therefore $2(k - j)$. Since in the minimal configuration there are n_k particles of charge k to the left of $p_{j,1}$, this gives a total contribution $\sum_{k=j+1}^{r-2} (k - j) n_k$. Apart from this, we encounter the situation where immediately to the left of $p_{j,1}$ we have a segment of the right-boundary particle. In such an instant we can perform L'_d , moving $p_{j,1}$ one step down. By construction of the minimal configuration, this occurs $\min(b - 1, r - j - 2)$ times. Finally, after having descended all the way down and having exchanged position with all particles of charge $> j$, $p_{j,1}$ is positioned immediately to the right of the left-boundary particle. It can then move up exactly $\min(a - 1, r - j - 2)$ times using move L'_u . Adding up all the contributions gives (3.20).

To see that (3.20) is unaltered by first having moved some (or all) particles of charge greater than j , consider a sequence of four points $X = \{(x_1, y_1), (x_2, y_2), (x_3, y_3), (x_4, y_4)\}$ connected by straight lines. First, let $y_1 > y_2 < y_3 > y_4$ and let $p_{j,1}$ be positioned immediately to the right of the sequence, i.e., the origin of $p_{j,1}$ is at (x_4, y_4) . Also, let the contour between the first two points not belong to the left-boundary particle. The total number of L -type steps $p_{j,1}$ can make is then

$$(y_3 - y_4 - j) + (y_3 - y_2 - j) + (y_1 - y_2 - j) = x_4 - x_1 - 3j$$

which is independent of the positions of the points (x_2, y_2) and (x_3, y_3) . Hence, carrying out any moves to X does not change the number of moves $p_{j,1}$ can make relative to X . If the contour between the first two points does belong to the left-boundary particle, this is changed to $x_4 - x_1 - 3j + r - 2 - \max(r - 2 - j, y_1)$, which is still independent of the relative positions of (x_2, y_2) and (x_3, y_3) . Second, let $y_1 < y_2 > y_3 < y_4$ and let $p_{j,1}$ be

positioned immediately, to the left of the sequence, i.e., the endpoint of $p_{j,1}$ is at (x_1, y_1) . Also, let the contour between the last two points not belong to the left-boundary particle. The total number of R -type steps $p_{j,1}$ can make is then

$$(y_2 - y_1 - j) + (y_2 - y_3 - j) + (y_4 - y_3 - j) = x_4 - x_1 - 3j$$

which is independent of the positions of the points (x_2, y_2) and (x_3, y_3) . Thanks to reversibility, the number of L -type moves $p_{j,1}$ can make relative to X is also $x_4 - x_3 - 3j$. If the contour between the last two points does belong to the right-boundary particle, this again changes by a term independent of the detailed positions of (x_2, y_2) and (x_3, y_3) . ■

Lemma 8. The maximal number of L -type moves $p_{j,l}$ can make is $k_{j,l-1}$, with $k_{j,l-1}$ the actual number of steps taken by $p_{j,l-1}$.

At last! We finally encounter the fermionic nature of our lattice gas.

Proof. Assume $p_{j,l-1}$ has made $k_{j,l-1}$ moves. Obviously, (before) the first $k_{j,l-1}$ moves, $p_{j,l}$ “sees” the same contour immediately to its left as $p_{j,l-1}$ did, when carrying out its leftward motion. Since $p_{j,l-1}$ and $p_{j,l}$ are identical particles, $p_{j,l}$ can thus carry out at least $k_{j,l-1}$ moves. Let $p_{j,l}$ indeed carry out $k_{j,l-1}$ moves. After that we encounter the situation of two pure particles of charge j , with the endpoint of the first being the origin of the next. The rightmost of the two can neither carry out L_u or L_d , since (in the notation of Definition 8) $y_1 = y_3$. ■

We note that the above two lemmas justify the chosen ordering of carrying out the leftward moves. First of all, by Lemma 8 it follows that we indeed have to move $p_{j,l-1}$ before $p_{j,l}$. Furthermore, we have to move $p_{k,m}$ before $p_{j,l}$, $k > j$, since the elementary moves only allow for leftward motion of pure particles; see Fig. 7. Finally, we have seen in the proof of Lemma 7 that the number of moves the particles of charge j can make is independent of the actual configuration of particles of charge $> j$.

Lemma 9. The contribution to the generating function \mathcal{C} of the particles of charge j is given by

$$\mathcal{C}_j = \begin{bmatrix} m_j + n_j \\ n_j \end{bmatrix} \tag{3.21}$$

Proof. From Lemmas 6–8 we get (dropping the subscripts j in the k -variables)

$$\mathcal{G}_j = \sum_{k_1=0}^{m_j} \sum_{k_2=0}^{k_1} \dots \sum_{k_{n_j}=0}^{k_{n_j}-1} q^{k_1+k_2+\dots+k_{n_j}} \quad (3.22)$$

We can (re)interpret this sum as the generating function of all partitions with largest part less than or equal to m_j and number of parts less than or equal to n_j . Thus we get (3.21); see, e.g., ref. 9.

Combining the above lemma with Lemma 1, we can state our second proposition as follows.

Proposition 2. The partition function of the one-dimensional Fermi gas is given by

$$Z(\mathbf{n}; a, b) = q^{E_{\min}} \prod_{j=1}^{r-3} \mathcal{G}_j = q^{E_{\min}} \prod_{j=1}^{r-3} \begin{bmatrix} m_j + n_j \\ n_j \end{bmatrix} \quad (3.23)$$

with E_{\min} given by (3.16) and m_j by (3.20).

To recast this result into a simpler form, we eliminate the n -variables in favor of the m -variables. To do so we use the simple formulas

$$\begin{aligned} -\min(p, q-1) + 2\min(p, q) - \min(p, q+1) &= \delta_{p, q}, & p, q-1 \geq 0 \\ -\min(p, q-1) + 2\min(p, q) &= p + \delta_{p, q}, & 0 \leq p \leq q+1 \end{aligned} \quad (3.24)$$

to get

$$-m_{j-1} + 2m_j - m_{j+1} = L\delta_{j,1} + \delta_{j,r-a-1} + \delta_{j,r-b-1} - 2n_j, \quad j = 1, \dots, r-3 \quad (3.25)$$

with $m_0 = m_{r-2} = 0$. To obtain the $j=1$ case of the above equation we made use of the completeness relation (3.8). Introducing the $(r-3)$ -dimensional vectors \mathbf{m} and \mathbf{e}_j with entries $(\mathbf{m})_j = m_j$ and $(\mathbf{e}_j)_k = \delta_{j,k}$, we can rewrite (3.25) as

$$\mathbf{n} = \frac{1}{2}(L\mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-b-1} - C\mathbf{m}) \quad (3.26)$$

Substituting this into Eq. (3.16) and (3.23), we arrive at the following simple result.

Proposition 3. The partition function of the Fermi gas of content \mathbf{n} reads

$$Z(\mathbf{n}; a, b) = q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \prod_{j=1}^{r-3} \left[\begin{array}{c} \frac{1}{2}(\mathcal{J} \mathbf{m} + L \mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-a-1})_j \\ m_j \end{array} \right] \quad (3.27)$$

with \mathbf{m} obtained through Eq. (3.26).

3.4. Computation of $\Xi_L(a, b)$

Having computed the partition function of our Fermi gas, it is only a trivial step to obtain the grand-canonical partition function $\Xi_L(a, b)$, defined in (3.6). In particular,

$$\Xi_L(a, b) = \sum_{\mathbf{n}} Z(\mathbf{n}; a, b) \quad (3.28)$$

Since our final result (3.27) for Z is entirely expressed through the m -variables, it is natural to also express the above sum over \mathbf{n} in terms of a sum over \mathbf{m} . From (3.20) and the fact that the occupation numbers n_j cannot be negative, we get

$$m_j = \min(a-1, r-j-2) + \min(b-1, r-j-2) + 2\mathbb{Z}_{\geq 0}, \quad j = 1, \dots, r-3 \quad (3.29)$$

Hence we obtain the grand-canonical partition function as

$$\begin{aligned} \Xi_L(a, b) &= \sum_{\mathbf{m}}^{(0)} q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \\ &\quad \times \prod_{j=1}^{r-3} \left[\begin{array}{c} \frac{1}{2}(\mathcal{J} \mathbf{m} + L \mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-a-1})_j \\ m_j \end{array} \right] \end{aligned} \quad (3.30)$$

where the (0) in the sum over \mathbf{m} denotes the restriction (3.29).

3.5. Computation of $X_L(a, b)$

To finally obtain the one-dimensional configuration sum $X_L(a, b)$, we have to carry out the sum (3.5), where we recall that $\Xi_L(a, b) := \Xi_L(a, b, 0, r)$.

To get the expression for $\Xi_L(a-\mu, b-\mu, 0, r-\mu)$, we have to make the substitutions $a \rightarrow a-\mu, b \rightarrow b-\mu$, and $r \rightarrow r-\mu$ in (3.30). This exactly gives back (3.30) apart from the fact that the restriction on the sum changes to

$$m_j = \begin{cases} \min(a-1, r-j-2) \\ \quad + \min(b-1, r-j-2) - 2\mu + 2Z \geq 0, & j = 1, \dots, r-\mu-3 \\ 0, & j = r-\mu-2, \dots, r-3 \end{cases} \tag{3.31}$$

Denoting this restriction as (μ) , we can write

$$X_L(a, b) = \sum_{\mu=0}^{\min(a,b)-1} \sum_{\mathbf{m}}^{(\mu)} q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \times \prod_{j=1}^{r-3} \left[\begin{matrix} \frac{1}{2}(\mathcal{J} \mathbf{m} + L \mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-a-1})_j \\ m_j \end{matrix} \right] \tag{3.32}$$

Combining the sum over \mathbf{m} restricted to (μ) and the sum over μ gives

$$X_L(a, b) = \sum'_{\mathbf{m}} q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \times \prod_{j=1}^{r-3} \left[\begin{matrix} \frac{1}{2}(\mathcal{J} \mathbf{m} + L \mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-a-1})_j \\ m_j \end{matrix} \right] \tag{3.33}$$

with the prime denoting yet another restriction,

$$m_j \equiv \min(a-1, r-j-2) + \min(b-1, r-j-2), \quad j = 1, \dots, r-3 \tag{3.34}$$

Unfortunately, we have not found an elegant way to prove this simplification and we defer it till the appendix.

To rewrite the above form of the restriction in the form conjectured in refs. 5 and 10 we note the identity

$$\begin{aligned} \min(p, q) \equiv & \delta_{p+1, q} + \delta_{p+3, q} + \delta_{p+5, q} + \dots \\ & + \delta_{1, q} + \delta_{3, q} + \delta_{5, q} + \dots \end{aligned} \tag{3.35}$$

for $p, q \geq 0$. Using this twice, once setting $p = a-1$ and $q = r-j-2$, and once setting $p = b-1$ and $q = r-j-2$, we get $m_j \equiv (\mathbf{Q}_{a,b})_j$, with $\mathbf{Q}_{a,b}$ given by (2.2).

We can thus conclude this section by formulating our main result as a theorem.

Theorem 1. For all $a = 1, \dots, r-1, b = 1, \dots, r-2,$ and $L - |a - b| \in 2\mathbb{Z}_{\geq 0},$ the one-dimensional configuration sum (1.5) is given by

$$\begin{aligned}
 X_L(a, b) = & \sum_{\mathbf{m} \in \mathbf{Q}_{a,b}} q^{(1/4)\mathbf{m}^T C \mathbf{m} - (1/2)m_{r-a-1}} \\
 & \times \prod_{j=1}^{r-3} \left[\frac{1}{2}(\mathcal{J} \mathbf{m} + L\mathbf{e}_1 + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-a-1})_j \right] \quad (3.36) \\
 & \qquad \qquad \qquad m_j
 \end{aligned}$$

where $\mathbf{Q}_{a,b}$ is given by (2.2).

4. BOSONIC SOLUTION OF THE ABF MODEL

In this section we recall the method for computing the sum (1.5) to obtain (up to a prefactor) the right-hand side of Melzer’s identities (2.1). This alternative approach to the sum (1.5) is the one originally taken by Andrews, Baxter, and Forrester⁽¹²⁾ and is given here mainly for reasons of completeness.

As a first step we introduce a function $Y_L(a, b)$ defined exactly as for $X_L(a, b)$ in (1.5), but with $\sigma_{L+1} = b - 2$ instead of $\sigma_{L+1} = b.$ We can then immediately infer the recurrence relations

$$X_L(a, b) = Y_{L-1}(a, b + 1) + q^{L/2} X_{L-1}(a, b - 1), \quad 1 \leq b \leq r - 2 \quad (4.1)$$

$$Y_L(a, b) = X_{L-1}(a, b - 1) + q^{L/2} Y_{L-1}(a, b + 1), \quad 2 \leq b \leq r - 1 \quad (4.2)$$

subject to the initial and boundary conditions

$$X_0(a, b) = Y_0(a, b) = \delta_{a,b} \quad (4.3)$$

$$X_L(a, 0) = Y_L(a, r) = 0 \quad (4.4)$$

To state the solution to these equations, we quote the following theorem established by Andrews, Baxter, and Forrester:⁽¹²⁾

Theorem 2. For $L \geq 0, 1 \leq a, b, c \leq r - 1, c = b \pm 1, L + a - b \equiv 0,$ let $X_L(a, b, c) = X_L(a, b)$ if $c = b + 1$ and $X_L(a, b, c) = Y_L(a, b)$ if $c = b - 1.$ Then

$X_L(a, b, c)$

$$= q^{(a-b)(a-c)/4} \sum_{j=-\infty}^{\infty} \left\{ q^{j(r(r-1)j+r(b+c-1)/2-(r-1)a)} \left[\begin{matrix} L \\ \frac{1}{2}(L+a-b)-rj \end{matrix} \right] \right. \\ \left. - q^{((r-1)j+(b+c-1)/2)(rj+a)} \left[\begin{matrix} L \\ \frac{1}{2}(L-a-b)-rj \end{matrix} \right] \right\} \quad (4.5)$$

To prove this, we note that (4.5) satisfies (4.1), thanks to

$$\begin{bmatrix} N \\ m \end{bmatrix} = \begin{bmatrix} N-1 \\ m-1 \end{bmatrix} + q^m \begin{bmatrix} N-1 \\ m \end{bmatrix} \quad (4.6)$$

Similarly, the proof that (4.5) satisfies (4.2) follows by application of

$$\begin{bmatrix} N \\ m \end{bmatrix} = \begin{bmatrix} N-1 \\ m \end{bmatrix} + q^{N-m} \begin{bmatrix} N-1 \\ m-1 \end{bmatrix} \quad (4.7)$$

To show that the initial condition (4.3) holds, note (1.3) as well as the range of a and b . This gives $j=0$ as the only nonvanishing term in the sum, and hence $X_0(a, b, c) = \delta_{a,b}$. Finally, $X_L(a, 0) = 0$ follows from (4.5) upon substitution of $b=0, c=1$ and making the change of variables $j \rightarrow -j$ in the first term within the curly braces. Analogously, $Y_L(a, r)$ follows from (4.5) upon substituting $b=r, c=r-1$ and making the change of variables $j \rightarrow -j-1$ in the first term within the braces. ■

To obtain the desired expression for $X_L(a, b)$, we set $c=b+1$ in (4.5), yielding

$$X_L(a, b) = f_{a,b}^{-1} \sum_{j=-\infty}^{\infty} \left\{ q^{j(r(r-1)j+rb-(r-1)a)} \left[\begin{matrix} L \\ \frac{1}{2}(L+a-b)-rj \end{matrix} \right] \right. \\ \left. - q^{((r-1)j+b)(rj+a)} \left[\begin{matrix} L \\ \frac{1}{2}(L-a-b)-rj \end{matrix} \right] \right\} \quad (4.8)$$

Combined with Theorem 1 at the end of Section 3, this proves Melzer's polynomial identities (2.1).

In conclusion to this section we make two remarks about the solution (4.8). First, the introduction of the auxiliary function $Y_L(a, b)$ could have been avoided, since from the definition (1.5) one can obtain recurrences that only involve the function $X_L(a, b)$. In particular,

$$X_L(a, b) = \begin{cases} q^{L/2} X_{L-1}(a, b-1) + q^{(L-1)/2} X_{L-1}(a, b+1) \\ \quad + (1 - q^{L-1}) X_{L-2}(a, b), & b = 1, \dots, r-3 \\ q^{L/2} X_{L-1}(a, b-1) + X_{L-2}(a, b), & b = r-2 \end{cases} \quad (4.9)$$

with the same conditions on $X_L(a, b)$ as in (4.3) and (4.4). The price to be paid for this is that, in order to show that (4.8) solves these relations, we need double application of (4.6) and (4.7). Interestingly, though, in terms of the fermionic left-hand side of (2.1), the recurrences (4.9) seem to be more natural; see e.g., ref. 11.

A second remark we wish to make is that like the fermionic result (2.1), also (4.8) has a nice interpretation in terms of restricted lattice paths. To see this, note that in order to obtain the generating function for the restricted lattice paths, we can first compute the generating function $G_L(\emptyset)$ of restricted lattice paths without the restriction $0 \leq y \leq r-2$. Since all paths that go below $y=0$ and above $y=r-2$ have now incorrectly been included, we have to subtract the generating function $G_L(\downarrow)$ of paths that somewhere go below $y=0$, as well as the generating function $G_L(\uparrow)$ of paths that somewhere go above $y=r-2$. However, we are again in error, since paths that go below $y=0$ as well as above $y=r-2$ have been subtracted twice. To correct this, we add $G_L(\downarrow, \uparrow)$ and $G_L(\uparrow, \downarrow)$, which are, respectively, the generating function of all paths that somewhere go above $y=r-2$ after having gone below $y=0$ and the generating function of all paths that somewhere go below $y=0$ after having gone above $y=r-2$. Again this is no good, and we keep continuing the process of adding and subtracting generating functions. In formula this reads

$$\begin{aligned}
 X_L(a, b) = \sum_{j \geq 0} \{ & G_L(\underbrace{\downarrow \uparrow \cdots \downarrow \uparrow}_{2j}) + G_L(\underbrace{\uparrow \downarrow \cdots \uparrow \downarrow}_{2j+2}) \\
 & - G_L(\underbrace{\downarrow \uparrow \cdots \uparrow \downarrow}_{2j+1}) - G_L(\underbrace{\uparrow \downarrow \cdots \downarrow \uparrow}_{2j+1}) \} \quad (4.10)
 \end{aligned}$$

with

$$G_L(\underbrace{\downarrow \uparrow \cdots \downarrow \uparrow}_{2j})$$

the generating function of all lattice paths that contain a sequence of extrema $\{(x_1, y_1), (x_2, y_2), \dots, (x_{2j}, y_{2j})\}$, with $x_j > x_k$ for $j > k$, $y_{2k-1} < 0$, and $y_{2k} > r-2$, and with the other generating functions defined similarly. Of course, since we consider paths of fixed, finite length, the above series only contains a finite number of nonzero terms. Computing the functions G_L , we obtain

$$\begin{aligned}
 G_L(\underbrace{\downarrow \uparrow \cdots \downarrow \uparrow}_{2j}) &= f_{a,b}^{-1} q^{j(r(r-1)j - rb + (r-1)a)} \left[\begin{matrix} L \\ \frac{1}{2}(L - a + b) - rj \end{matrix} \right] \\
 G_L(\underbrace{\uparrow \downarrow \cdots \uparrow \downarrow}_{2j}) &= f_{a,b}^{-1} q^{j(r(r-1)j + rb - (r-1)a)} \left[\begin{matrix} L \\ \frac{1}{2}(L + a - b) - rj \end{matrix} \right] \\
 G_L(\underbrace{\downarrow \uparrow \cdots \downarrow \uparrow}_{2j+1}) &= f_{a,b}^{-1} q^{((r-1)j + b)(rj + a)} \left[\begin{matrix} L \\ \frac{1}{2}(L - a - b) - rj \end{matrix} \right] \\
 G_L(\underbrace{\uparrow \downarrow \cdots \uparrow \downarrow}_{2j+1}) &= f_{a,b}^{-1} q^{((r-1)(j+1) - b)(r(j+1) + a)}
 \end{aligned} \tag{4.11}$$

$$\times \left[\begin{matrix} L \\ \frac{1}{2}(L + a + b) - r(j+1) \end{matrix} \right] \tag{4.12}$$

for all $j \geq 0$. Substitution in (4.10) correctly reproduces the expression (4.8). We remark that the above method for computing $X_L(a, b)$ is merely a rewording of the *sieving* technique developed by Andrews in the context of partition theory; see, e.g., ref. 9. For the details of the calculation leading to (4.11) we refer the reader to ref. 9, Chapter 9, and ref. 24.

5. SUMMARY AND DISCUSSION

In this paper we have computed, using the combinatorial technique developed in part I, all one-dimensional configuration sums of the $(r - 1)$ -state ABF model. In contrast to the earlier results of Andrews, Baxter, and Forrester, our expressions are of so-called fermionic type, and provide a new proof of polynomial identities conjectured by Melzer. In the limit of an infinitely large lattice, these identities imply the fermionic expressions for the $\chi_{b,a}^{(r-1,r)}(q)$ Virasoro characters as conjectured by the Stony Brook group. Using the Andrews–Bailey construction, we also proved fermionic expressions for several nonunitary minimal Virasoro characters.

In conclusion to this paper we make a few comments. First, motivated by the ground-breaking papers of the Stony Brook group,^(5, 19, 20, 25, 26) a vast literature has arisen containing numerous claims for identities of the Rogers–Ramanujan type.^(10, 27–34) We expect that our fermionic method for computing generating functions of restricted lattice paths can be applied to obtain proofs of several of these conjectures. Other recently developed approaches toward either proof or increased understanding of Fermi–Bose character identities can be found, e.g., in refs. 6, 7, 11, and 35–47.

A second remark is that in the $q = 1$ limit, Melzer's identities reduce to identities for the number of L -step paths on the A_{r-1} Dynkin diagram, with fixed initial and final positions. Viewed in this way, it turns out that Melzer's identities are in fact a special one-dimensional case of polynomial identities for q -deformed path counting on arbitrary d -dimensional cuboids. In the limit of infinitely long paths, these "cuboid" identities decouple into products of Virasoro character identities. The simplest example beyond Melzer's case is the q -deformation of a path-counting formula on a "railroad" of length $r - 3$, reading

$$\begin{aligned}
 f_{a,b} & \sum_{\mathbf{m} \equiv \mathbf{Q}_{a,b}} q^{(1/4)\mathbf{m}^T \mathbf{C} \mathbf{m} - (1/2)m_{r-a-1}} \left[\begin{matrix} L \\ m_1 \end{matrix} \right] \prod_{j=2}^{r-3} \left[\begin{matrix} \frac{1}{2}(\mathcal{J} \mathbf{m} + \mathbf{e}_{r-a-1} + \mathbf{e}_{r-b-1})_j \\ m_j \end{matrix} \right] \\
 & = \sum_{j=-\infty}^{\infty} \left\{ q^{j(r(r-1)j+rb-(r-1)a)} \left[\begin{matrix} L \\ 2(r-1)j+b-a \end{matrix} \right]_2 \right. \\
 & \quad \left. - q^{((r-1)j+b)(rj+a)} \left[\begin{matrix} L \\ 2(r-1)j+b+a \end{matrix} \right]_2 \right\} \tag{5.1}
 \end{aligned}$$

for $a, b = 1, \dots, r - 2$ and $L \geq 0$. Here $\left[\begin{matrix} N \\ m \end{matrix} \right]_2$ are q -deformed trinomial coefficients defined as⁽⁴⁸⁾

$$\left[\begin{matrix} N \\ m \end{matrix} \right]_2 = \sum_{k \geq 0} q^{k(k+m)} \left[\begin{matrix} N \\ k \end{matrix} \right] \left[\begin{matrix} N-k \\ k+m \end{matrix} \right] \tag{5.2}$$

A discussion for the case of arbitrary cuboids will be presented elsewhere.⁽⁴⁹⁾

A final remark is that the result (3.36) proven in this paper has nice partition-theoretic interpretations. One follows from the work of Andrews *et al.*,⁽²⁴⁾ stating that the one-dimensional configuration sum $X_L(a, b)$ is the generating function of all partitions into at most $\frac{1}{2}(L + a - b)$ parts, each part $\leq \frac{1}{2}(L - a + b)$, such that the hook differences on the $(1 - b)$ th diagonal are $\geq b - a + 1$ and those on the $(r - b - 2)$ th diagonal are $\leq b - a$. Another interpretation follows by viewing a restricted lattice path with total energy E as a partition of $2E = \lambda_1 + \lambda_2 + \dots + \lambda_M$, with λ_j the j th x -position counted from the right, where the path has no extremum. With this map from paths to partitions, $X_L(a, b; q^2)$ is the generating function of partitions into parts $\lambda_1, \lambda_2, \dots, \lambda_M$, with $\lambda_M < \dots < \lambda_2 < \lambda_1 \leq L$, and

$$1 - b \leq u_j - d_j \leq r - b - 2, \quad \forall j = 1, \dots, M \tag{5.3}$$

$$u_M - d_M = a - b, a - b - 1$$

Here u_j is the number of parts $\lambda_k \equiv a - b + k$ for $k \leq j$ and d_j is the number of parts $\lambda_k \not\equiv a - b + k$ for $k \leq j$.

APPENDIX A. PROOF OF EQUIVALENCE OF (3.32) AND (3.33)

In this appendix we prove that (3.32) can be simplified to yield the final result (3.33) for the one-dimensional configuration sums.

The problem with this step is that the following statement turns out to be false:

$$\sum_{\mu=0}^{\min(a,b)-1} \sum_{\mathbf{m}}^{(\mu)} = \sum_{\mathbf{m}}' \quad (\text{A.1})$$

where (μ) denotes the restriction (3.31) and the prime the restriction (3.34). In particular, the number of vectors \mathbf{m} which are in accordance with the restriction (3.34) exceed the number of vectors \mathbf{m} in accordance with the restriction (3.31) summed over μ . What we will show now is that each of the additional \mathbf{m} 's allowed by the sum on the right-hand side gives a vanishing contribution.

Let us assume that $\min(a, b) = M + 1$, so that we have $M + 1$ terms in the sum over μ . Let us further define S_μ as the set of \mathbf{m} -vectors allowed by the restriction (3.31). In other words, $\mathbf{m} \in S_\mu$

$$\begin{cases} m_1, \dots, m_{r-M-3} \geq M - \mu \\ m_{r-M+k-3} \geq M - \mu - k + 1, & k = 1, \dots, M - \mu \\ m_{r-\mu-2} = \dots = m_{r-3} = 0 \\ m_j \equiv \min(a-1, r-j-2) + \min(b-1, r-j-2), & j = 1, \dots, r-3 \end{cases} \quad (\text{A.2})$$

Note that $S_\mu \cap S_\nu = \emptyset$ for $\mu \neq \nu$. We now use that $[\binom{N}{m}]$ is nonvanishing for $0 \leq m \leq N$ only. From this and the summand of (3.32), we infer

$$0 \leq m_j \leq \frac{1}{2}(m_{j-1} + m_{j+1} + \delta_{r-a-1, j} + \delta_{r-b-1, j}), \quad j = 2, \dots, r-3 \quad (\text{A.3})$$

with $m_{r-2} = 0$. From this one immediately sees that

$$m_j \geq m_k \quad \text{for } j < k \quad (\text{A.4})$$

However, interestingly enough, this condition is not yet good enough for our purposes. Instead we need to use the fact that $\min(a, b) = M + 1$. This combined with (A.3) gives rise to the following ordering for all $j = r - M - 1, \dots, r - 3$:

$$\text{if } m_j \geq 1 \Rightarrow m_{j-k} \geq k + 1, \quad k = 1, \dots, M - r + j + 2 \quad (\text{A.5})$$

in addition to (A.4).

In the following we use the notation $A \cup B = C$ to mean $A \cup B \subseteq C$ and

$$\sum_{\mathbf{m} \in A} \sum_{\mathbf{m} \in B} f(\mathbf{m}) = \sum_{\mathbf{m} \in C} f(\mathbf{m}) \tag{A.6}$$

with $f(\mathbf{m})$ the summand in (3.32).

We now proceed by induction. We set $T_n = T_{n-1} \cup S_{M-n}$, with $T_0 = S_M$, and claim that T_n is given by

$$\begin{cases} m_1, \dots, m_{r-M+n-3} \geq 0 \\ m_{r-M+n-2} = \dots = m_{r-3} = 0 \\ m_j \equiv \min(a-1, r-j-2) + \min(b-1, r-j-2) \end{cases} \tag{A.7}$$

For $n=0$ this is correct by construction. To show the induction step, set $\mu = M - n - 1$ in (A.2). This yields

$$\begin{cases} m_1, \dots, m_{r-M-3} \geq n+1 \\ m_{r-M+k-3} \geq n-k+2, \quad k=1, \dots, n+1 \\ m_{r-M+n-1} = \dots = m_{r-3} = 0 \\ m_j \equiv \min(a-1, r-j-2) + \min(b-1, r-j-2) \end{cases} \tag{A.8}$$

Using the conditions (A.4) and (A.5) with $j=r-M+n-2$, we can combine the above two equations to find that T_{n+1} is given by (A.7) with n replaced by $n+1$. This proves our claim (A.7) and we obtain the desired expression for T_M by setting $n=M$ in (A.7). This indeed gives the restriction (3.34) we set out to prove. ■

ACKNOWLEDGMENTS

I wish to thank Alexander Berkovich, Omar Foda, Peter Forrester, and Barry McCoy for many interesting and helpful discussions. This work was supported by the Australian Research Council. I thank a referee for pointing out some misprints in an earlier version of this manuscript.

Note Added. After completing this manuscript I received a preprint by A. Schilling⁽⁵⁰⁾ in which (2.1) as well as (2.4) are proven as special cases of polynomial identities for finitized branching functions of the cosets $\widehat{su}(2)_M \times \widehat{su}(2)_N / \widehat{su}(2)_{N+M}$. The result of ref. 50 is based on recursion techniques and generalizes the work of Berkovich.⁽¹¹⁾

REFERENCES

1. L. J. Rogers, *Proc. Lond. Math. Soc.* **25**:318 (1894).
2. L. J. Rogers, *Proc. Camb. Phil. Soc.* **19**:211 (1919).
3. S. Ramanujan, *Proc. Camb. Phil. Soc.* **19**:214 (1919).
4. A. Rocha-Caridi, In *Vertex Operators in Mathematics and Physics*, J. Lepowsky, S. Mandelstam, and I. M. Singer, eds. (Springer, Berlin, 1985).
5. R. Kedem, T. R. Klassen, B. M. McCoy, and E. Melzer, *Phys. Lett.* **307B**:68 (1993).
6. O. Foda and Y.-H. Quano, Virasoro character identities from the Andrews–Bailey construction, Preprint University of Melbourne No. 26–94 [hep-th/9408086].
7. A. Berkovich and B. M. McCoy, Continued fractions and fermionic representations for characters of $M(p, p')$ minimal models, Preprint BONN-TH-94-28, ITPSB 94-060 [hep-th/9412030]; *Lett. Math. Phys.*, to appear.
8. I. J. Schur, *Sitzungsber. Preuss. Akad. Wiss. Phys.-Math. Kl.* **1917**:302.
9. G. E. Andrews, *The Theory of Partitions* (Addison-Wesley, Reading, Massachusetts, 1976).
10. E. Melzer, *Int. J. Mod. Phys. A* **9**:1115 (1994).
11. A. Berkovich, *Nucl. Phys. B* **431**:315 (1994).
12. G. E. Andrews, R. J. Baxter, and P. J. Forrester, *J. Stat. Phys.* **35**:193 (1984).
13. R. J. Baxter, *J. Stat. Phys.* **26**:427 (1981).
14. W. N. Bailey, *Proc. Lond. Math. Soc.* (2) **49**:421 (1947).
15. W. N. Bailey, *Proc. Lond. Math. Soc.* (2) **50**:1 (1949).
16. G. E. Andrews, *Pac. J. Math.* **114**:267 (1984).
17. S. O. Warnaar, *J. Stat. Phys.* **82**:657 (1996).
18. A. Berkovich, Private communication.
19. R. Kedem, T. R. Klassen, B. M. McCoy, and E. Melzer, *Phys. Lett.* **304B**:263 (1993); **307B**:68 (1993).
20. S. Dasmahapatra, R. Kedem, T. R. Klassen, B. M. McCoy, and E. Melzer, *Int. J. Mod. Phys. B* **7**:3617 (1993).
21. D. Bressoud, In *Lecture Notes in Mathematics*, Vol. 1395 (Springer-Verlag, Berlin, 1987), p. 140.
22. B. Gordon, *Amer. J. Math.* **83**:393 (1961).
23. G. E. Andrews, *Proc. Natl. Acad. Sci. USA* **74**:4082 (1974).
24. G. E. Andrews, R. J. Baxter, D. M. Bressoud, W. J. Burge, P. J. Forrester, and G. Viennot, *Eur. J. Comb.* **8**:341 (1987).
25. R. Kedem and B. M. McCoy, *J. Stat. Phys.* **71**:883 (1993).
26. S. Dasmahapatra, R. Kedem, B. M. McCoy, and E. Melzer, *J. Stat. Phys.* **74**:239 (1994).
27. A. Kuniba, T. Nakanishi, and J. Suzuki, *Mod. Phys. Lett. A* **8**:1649 (1993).
28. S. Dasmahapatra, String hypothesis and characters of cosets CFT's, Preprint ICTP IC/93/91 [hep-th/9305024].
29. S. Dasmahapatra, *Int. J. Mod. Phys. A* **10**:875 (1995).
30. E. Melzer, *Lett. Math. Phys.* **31**:233 (1994).
31. S. O. Warnaar and P. A. Pearce, *Int. J. Mod. Phys. A* **11**:291 (1996).
32. E. Melzer, Supersymmetric analogs of the Gordon–Andrews identities, and related TBA systems, Preprint TAUP 2211-94 [hep-th/9412154].
33. D. Gepner, *Phys. Lett.* **348B**:377 (1995).
34. E. Bayer and D. Gepner, Fermionic sum representations for the Virasoro characters of the unitary superconformal minimal models, Preprint [hep-th/9502118].
35. P. Bouwknegt, A. W. W. Ludwig, and K. Schoutens, *Phys. Lett.* **338B**:448 (1994).

36. P. Bouwknegt, A. W. W. Ludwig, and K. Schoutens, Spinon basis for higher level $Su(2)$ WZW models, Preprint USC-94/20 [hep-th/9412108].
37. P. Bouwknegt, A. W. W. Ludwig, and K. Schoutens, Affine and Yangian symmetries in $SU(2)_1$ conformal field theory, Preprint USC-94/21 [hep-th/9412199].
38. O. Foda and Y.-H. Quano, *Int. J. Mod. Phys.* **10**:2291 (1995).
39. S. O. Warnaar and P. A. Pearce, *J. Phys. A: Math. Gen.* **27**:L891 (1994).
40. G. Georgiev, Combinatorial construction of modules for infinite-dimensional Lie algebras, I. Principal subspace, Preprint Rutgers University [hep-th/9412054].
41. G. Georgiev, Combinatorial construction of modules for infinite-dimensional Lie algebras, II. Parafemionic space, Preprint Rutgers University [q-alg/9504024].
42. O. Foda and S. O. Warnaar, *Lett. Math. Phys.* **36**:145 (1996).
43. A. Nakayashiki and Y. Yamada, Crystallizing the spinon basis, Preprint Kyushu University [hep-th/9504052], *Commun. Math. Phys.*, to appear.
44. A. Nakayashiki and Y. Yamada, *Int. J. Mod. Phys. A* **11**:395 (1996).
45. A. Berkovich, B. M. McCoy, and W. P. Orrick, Polynomial identities, indices, and duality for the $N=1$ superconformal model $SM(2.4\nu)$, Preprint BONN-TH-95-12, ITPSB 95-18 [hep-th/9507072].
46. O. Foda, M. Okado, and S. O. Warnaar, A proof of polynomial identities of type $\widehat{sl(n)}_1 \otimes \widehat{sl(n)}_1 / \widehat{sl(n)}_2$, Preprint University of Melbourne [q-alg/9507014], *J. Math. Phys.*, to appear.
47. T. Tarakawa, T. Nakanishi, K. Oshima, and A. Tsuchiya, Spectral decomposition of path space in solvable lattice model, Preprint Nagoya University [q-alg/9507025].
48. G. E. Andrews and R. J. Baxter, *J. Stat. Phys.* **47**:297 (1987).
49. S. O. Warnaar, Path counting on hypercuboids and Rogers–Ramanujan type identities, In preparation.
50. A. Schilling, Polynomial fermionic forms for the branching functions of the rational coset conformal field theories $\widehat{su(2)}_M \times \widehat{su(2)}_N / \widehat{su(2)}_{N+M}$, Preprint ITPSB [hep-th/9508050], *Nucl. Phys. B*, to appear.